

The unipotent Albanese map and Selmer varieties for curves

Minhyong Kim

February 2, 2008

Abstract

We study the unipotent Albanese map that associates the torsor of paths for p -adic fundamental groups to a point on a hyperbolic curve. It is shown that the map is very transcendental in nature, while standard conjectures about the structure of mixed motives provide control over the image of the map. As a consequence, conjectures of ‘Birch and Swinnerton-Dyer type’ are connected to finiteness theorems of Faltings-Siegel type.

Dedicated to the memory of my teacher Serge Lang.

In a letter to Faltings [15] dated June, 1983, Grothendieck proposed several striking conjectural connections between the arithmetic geometry of ‘anabelian schemes’ and their fundamental groups, among which one finds issues of considerable interest to classical Diophantine geometers. Here we will trouble the reader with a careful formulation of just one of them. Let F be a number field and

$$f : X \rightarrow \mathrm{Spec}(F)$$

a smooth, compact, hyperbolic curve over F . After the choice of an algebraic closure

$$y : \mathrm{Spec}(\bar{F}) \rightarrow \mathrm{Spec}(F)$$

and a base point

$$x : \mathrm{Spec}(\bar{F}) \rightarrow X$$

such that $f(x) = y$, we get an exact sequence of fundamental groups:

$$0 \rightarrow \hat{\pi}_1(\bar{X}, x) \rightarrow \hat{\pi}_1(X, x) \xrightarrow{f_*} \Gamma \rightarrow 0,$$

where $\Gamma = \mathrm{Gal}(\bar{F}/F)$ is the Galois group of $\bar{F}$ over F and $\bar{X} = X \otimes \bar{F}$ is the base change of X to $\bar{F}$. Now, suppose we are given a section $s \in X(F)$ of the map $X \rightarrow \mathrm{Spec}(F)$, i.e., an F -rational point of X . This induces a map of fundamental groups

$$s_* : \Gamma \rightarrow \hat{\pi}_1(X, x')$$

where $x' = s(y)$. Choosing an étale path p from x' to x determines an isomorphism

$$c_p : \hat{\pi}_1(X, x') \simeq \hat{\pi}_1(X, x), \quad l \mapsto p \circ l \circ p^{-1},$$

which is independent of p up to conjugacy. On the other hand, p maps to an element $\gamma \in \Gamma$, and it is straightforward to check that $c_p \circ s_* \circ c_{\gamma^{-1}}$ is a continuous splitting of $f_* : \hat{\pi}_1(X, x) \rightarrow \Gamma$. This splitting is well-defined up to the equivalence relation $\sim$ given by conjugacy of sections, where $\gamma \in \hat{\pi}_1(X, x)$ acts on a section s as $s \mapsto c_\gamma \circ s \circ c_{f_*(\gamma)^{-1}}$. We get thereby a map

$$s \mapsto [c_p \circ s_* \circ c_{\gamma^{-1}}]$$

from $X(F)$ to the set $Split(X)/\sim$ of splittings $Split(X)$ of the exact sequence modulo the equivalence relation $\sim$ given by conjugation. Grothendieck's *section conjecture* states that this map is a bijection:

$$X(F) \simeq Split(X)/\sim$$

It seems that during the initial period of consideration, there was an expectation that the section conjecture would ‘directly imply’ the Mordell conjecture. At present the status of such an implication is unclear. Nevertheless, this fact does not diminish the conceptual importance of the section conjecture and its potential for broad ramifications in Diophantine geometry. In the long run, one can hope that establishing the correct link between Diophantine geometry and homotopy theory will provide us with the framework for a deeper understanding of Diophantine finiteness, especially in relation to the analytic phenomenon of hyperbolicity.

In this paper, we wish to make some preliminary comments on the fundamental groups/Diophantine geometry connection from a somewhat different perspective which, needless to say, does not approach the depth of the section conjecture. In fact, in our investigation, the main tool is the *motivic fundamental group*, especially p -adic realizations, rather than the pro-finite fundamental group. Nevertheless, what emerges is a (murky) picture containing at least a few intriguing points of mystery that are rather surprising in view of the relative poverty of unipotent completions. That is to say, one is surprised by the Diophantine depth of an invariant that is so close to being linear. It is the author's belief that the *Selmer varieties* arising in this context, generalizing $\mathbb{Q}_p$ -Selmer groups of abelian varieties, are objects of central interest. Developing the formalism for a systematic study of Selmer varieties is likely to be crucial for continuing research along the lines suggested in this paper. However, the non-abelian nature of the construction presents a formidable collection of obstacles that are at present beyond the author's power to surmount. In spite of this, it is hoped that even a partial resolution of the problems can point us eventually towards the full algebraic completion of the fundamental group, bringing a sort of ‘motivic Simpson theory’ to bear upon the study of Diophantine sets.

We proceed then to a brief summary of the notions to be discussed and a statement of the results, omitting precise definitions for the purposes of this introduction. Although it will be clear that at least part of the formalism is more general, we will focus our attention on curves in this paper, right at the outset. So we let F be a number field and X/F a smooth hyperbolic curve, possibly non-compact. Let R be the ring of S -integers in F for some finite set S of primes. We assume that we are given a smooth model

$$\begin{array}{ccc} X & \hookrightarrow & \mathcal{X} \\ \downarrow & & \downarrow \\ \mathrm{Spec}(F) & \hookrightarrow & \mathrm{Spec}(R) \end{array}$$

of X and a compactification $\mathcal{X} \hookrightarrow \mathcal{X}'$ relative to R where $\mathcal{X}'$ and the complement $\mathcal{D}$ of $\mathcal{X}$ in $\mathcal{X}'$ are also smooth over R . The Diophantine set of interest in this situation is $\mathcal{X}(R)$, the S -integral points of $\mathcal{X}$. The theorems of Siegel and Faltings say that this set is finite. After setting up the preliminary formalism, our goal will be to investigate these theorems from a π_1 -viewpoint.

As already mentioned, the π_1 we will be focusing on in this paper is a part of the *motivic* π_1 defined by Deligne [9]. In spite of the terminology, we do not rely on any general theory of motives in our discussion of the fundamental group. Rather, there will always be specific realizations related by the standard collection of comparison maps. Nevertheless, it is probably worth emphasizing that the main idea underlying our approach is that of a *motivic unipotent Albanese map*, of which the one defined by Hain [16] is the Hodge realization. That is, whenever a point $b \in \mathcal{X}(R)$ is chosen as a basepoint, a unipotent motivic fundamental group $U^{mot} := \pi_1^{mot}(X, b)$ as well as motivic torsors $P^{mot}(y) := \pi_1^{mot}(X; b, y)$ of paths associated to other points y should be determined. The idea is that if a suitable classifying space D^{mot} for such torsors were to be constructed, the Albanese map would merely associate to a point y , the class

$$[P^{mot}(y)] \in D^{mot}.$$

Coming back to the concrete consideration of realizations, the analytic part of the machinery we need comes from the *De Rham fundamental group* of $X_v := X \otimes F_v$ that depends only on the local

arithmetic geometry. Here, v is an archimedean valuation of F not contained in S . The definition of this fundamental group requires the use of the category $\text{Un}(X_v)$ of unipotent vector bundles with connections on X_v . Given any base point $b \in \mathcal{X}(R_v)$, we get a fiber functor $e_b : \text{Un}(X) \rightarrow \text{Vect}_{F_v}$ to the category of vector spaces over F_v , and

$$U^{DR} := \pi_{1,DR}(X_v) := \text{Aut}^\otimes(e_b)$$

in a rather obvious sense as functors on affine F_v -schemes. The functor $\pi_{1,DR}(X_v)$ ends up being representable by a pro-unipotent pro-algebraic group. Natural quotients

$$U_n^{DR} := Z^n \backslash U^{DR}$$

via the descending central series of U^{DR} correspond to restricting e_b to the subcategory generated by bundles having index of unipotency $\leq n$. These quotients are unipotent algebraic groups over F_v . U^{DR} is endowed with a decreasing *Hodge filtration* by subgroups:

$$U^{DR} \supset \dots \supset F^n U^{DR} \supset F^{n+1} U^{DR} \supset \dots \supset F^0 U^{DR}$$

coming from a filtration of the coordinate ring by ideals. There is a comparison isomorphism

$$U^{DR} \simeq U^{cr} \otimes_K F_v = \pi_{1,cr}(Y_v, \bar{b}) \otimes_K F_v$$

(where K is the maximal absolutely unramified subfield of F_v) with the crystalline fundamental group of the special fiber, defined using unipotent over-convergent iso-crystals. The utility of this is that the Frobenius of the special fiber comes to act naturally on the De Rham fundamental group. Of crucial importance for us is the consideration of De Rham ‘path spaces’:

$$P^{DR}(x) := \pi_{1,DR}(X_v; x, b) := \text{Isom}^\otimes(e_b, e_x)$$

consisting of isomorphisms from the fiber functor e_b to e_x , $x \in X_v(R_v)$. These are pro-algebraic varieties also endowed with Hodge filtrations and crystalline Frobenii that are compatible with their structure as right torsors for U^{DR} , and the point is to study how the data vary with x . In fact, they turn out to be classified by a natural ‘period space’

$$U^{DR}/F^0 U^{DR}$$

giving us a higher De Rham unipotent Albanese map

$$j^{DR} : \mathcal{X}(R_v) \rightarrow U^{DR}/F^0 U^{DR}$$

that maps x to the class of $P^{DR}(x)$. By passing to (finite-dimensional) quotients U_n^{DR} , we also have finite-level versions

$$j_n^{DR} : \mathcal{X}(R_v) \rightarrow U_n^{DR}/F^0 U_n^{DR}$$

that fit into a compatible tower

$$\begin{array}{ccc} \parallel & & \downarrow \\ \mathcal{X}(R_v) & \rightarrow & U_{n+1}^{DR}/F^0 \\ \parallel & & \downarrow \\ \mathcal{X}(R_v) & \rightarrow & U_n^{DR}/F^0 \\ \parallel & & \downarrow \\ \vdots & \vdots & \vdots \\ \parallel & & \downarrow \\ \mathcal{X}(R_v) & \rightarrow & U_2^{DR}/F^0 \end{array}$$

At the very bottom, $U_2^{DR} = H_1^{DR}(X_v) := (H_{DR}^1(X_v))^*$ and j_2^{DR} is nothing but the logarithm of the usual Albanese map with respect to the basepoint b .

A comparison with the situation over $\mathbb{C}$ easily yields the following:

Theorem 1 *For each $n \geq 2$, the image of j_n^{DR} is Zariski dense.*

This statement can be interpreted as linear-independence for multiple polylogarithms of higher genus. It has been pointed out by a referee that this theorem is also proved by Faltings in a preprint [10]. In fact, both Faltings and Akio tamagawa had indicated to the author earlier the possibility of proving the denseness using trascendental methods in the genus zero case.

Unfortunately, this simple theorem is the only concrete result to be reported on in this paper, the remaining parts being an extended commentary on what else might be expected when allowed considerable optimism. Nevertheless, we proceed to summarize here our observations.

The construction described thus far will have to be compared with one involving pro-unipotent étale fundamental groups that are associated to the situation in both local and global settings. That is, after a choice $\bar{F}_v$ of an algebraic closure for F_v , we can associate to the base-change $\bar{X}_v := X_v \otimes \bar{F}_v$ the pro-unipotent étale fundamental group

$$U^{et} := \pi_{1,\text{ét}}(\bar{X}_v, b)$$

that classifies unipotent lisse $\mathbb{Q}_p$ -sheaves on $\bar{X}_v$. Furthermore, to each x , we can also associate the space of unipotent étale paths $P^{et} = \pi_{1,\text{ét}}(\bar{X}_v; x, b)$ which is a torsor for U^{et} . Both carry actions of $G_v := \text{Gal}(\bar{F}_v/F_v)$ and the torsor structure is compatible with the action. The torsors that are associated in this way to integral points $x \in \mathcal{X}(R_v)$ have the additional property of being trivialized over a ring B_{cr} of p -adic periods. We classify these torsors using a restricted Galois cohomology set $H_f^1(G_v, U^{et})$. A classifying map defined exactly analogously to the De Rham setting provides a local étale Albanese map

$$j_{loc}^{et} : \mathcal{X}(R_v) \rightarrow H_f^1(G_v, U^{et})$$

as well as finite-level versions

$$(j_{loc}^{et})_n : \mathcal{X}(R_v) \rightarrow H_f^1(G_v, U_n^{et}).$$

The connection to j^{DR} goes through a non-abelian extension of Fontaine's Dieudonné functor, interpreted as an algebraic map

$$D : H_f^1(G_v, U_n^{et}) \rightarrow U_n^{DR}/F^0$$

that fits into a commutative diagram

$$\begin{array}{ccc} \mathcal{X}(R_v) & \rightarrow & U_n^{DR}/F^0 \\ & \searrow & \uparrow \\ & & H_f^1(G_v, U_n^{et}). \end{array}$$

The study of global points comes into the picture when we take the base-point b itself from $\mathcal{X}(R)$, the set of global integral points, and consider the global $\mathbb{Q}_p$ pro-unipotent étale fundamental group $\pi_{1,\text{ét}}(\bar{X}, b)$ with the action of $\Gamma = \text{Gal}(\bar{F}/F)$. A choice of an embedding $\bar{F} \hookrightarrow \bar{F}_v$ determines an inclusion $G_v \hookrightarrow \Gamma$. Other than the consideration of the Galois action, we have an isomorphism of pro-algebraic groups $\pi_{1,\text{ét}}(\bar{X}, b) \simeq \pi_{1,\text{ét}}(\bar{X}_v, b)$, so we will allow ourself a minimal abuse of notation and denote the global fundamental group also by U^{et} . Any other global point y then determines a torsor which is denoted $P^{et}(y)$ (again introducing a bit of confusion with the local object). We denote by T the set of primes consisting of S together with all primes dividing the residue characteristic of v . The action of G on U^{et} factors through G_T , the Galois group of the maximal subfield of $\bar{F}$ unramified over the primes in T . Of crucial importance for our purposes is the non-abelian cohomology set

$$H^1(G_T, U^{et})$$

and a natural subset

$$H_f^1(G_T, U^{et}) \subset H^1(G_T, U^{et})$$

defined by ‘Selmer conditions’ at the primes in T (which need not be too precisely determined in most situations). All the cohomology sets thus far discussed can be interpreted as the points of a pro-algebraic variety (over $\mathbb{Q}_p$), and $H_f^1(G_T, U^{et})$ is the Selmer variety occurring in the title of this paper. The finite-level versions $H_f^1(G_T, U_n^{et})$ and $H_f^1(G_v, U_n^{et})$ are finite-dimensional algebraic varieties. In this interpretation, the restriction map and the Dieudonné functor become algebraic maps of $\mathbb{Q}_p$ -schemes, the target of the latter being the Weil restriction $Res_{\mathbb{Q}_p}^{F_v}(U^{DR}/F^0)$, which of course has the property that

$$Res_{\mathbb{Q}_p}^{F_v}(U^{DR}/F^0)(\mathbb{Q}_p) = (U^{DR}/F^0)(F_v).$$

We have thus described the fundamental diagram

$$\begin{array}{ccccc} \mathcal{X}(R) & \rightarrow & \mathcal{X}(R_v) & \rightarrow & U_n^{DR}/F^0 \\ \downarrow & & \downarrow & \nearrow & \\ H_f^1(G_T, U_n^{et}) & \rightarrow & H_f^1(G_v, U_n^{et}) & & \end{array}$$

that provides for us the link between Diophantine geometry and the theory of fundamental groups. The reader familiar with the method of Chabauty [4] will recognize here a non-abelian lift of the diagram

$$\begin{array}{ccccc} \mathcal{X}(R) & \rightarrow & \mathcal{X}(R_v) & \rightarrow & \text{Lie}(J) \otimes F_v \\ \downarrow & & \downarrow & \nearrow & \\ J(R) \otimes \mathbb{Q}_p & \rightarrow & J(R_v) \otimes \mathbb{Q}_p & & \end{array}$$

where J is the Jacobian of $\mathcal{X}$, which is essentially the case $n = 2$, that is, other than the replacement of the Mordell-Weil group by the Selmer group. The point is that the image of $\mathcal{X}(R)$ inside U_n^{DR}/F^0 thus ends up being contained inside the image of $H_f^1(G_T, U_n^{et})$. The desired relation to Diophantine finiteness is expressed by the

Conjecture 1 *Let $F = \mathbb{Q}$ and X be hyperbolic. Then*

$$\dim(H_f^1(G_T, U_n^{et})) < \dim(U_n^{DR}/F^0)$$

for some n (possibly very large).

Whenever one can find an n for which this inequality of dimensions is verified, finiteness of $\mathcal{X}(R)$ follows, exactly as in Chabauty’s argument. The only difference is that the analytic functions that occur in the proof are p -adic *iterated integrals* [13] rather than abelian integrals [7].

Unfortunately, this conjecture can be proved at present only when the genus of X is zero [22] and some special genus one situations (with a slight modification of the Selmer variety). However, there is perhaps some interest in the tight connection we establish between the desired statement and various conjectures of ‘Birch and Swinnerton-dyer type.’ For example, our conjecture is implied by any one of:

- (1) a certain fragment of the Bloch-Kato conjecture [3];
- (2) the Fontaine-Mazur conjecture [12];
- (3) Jannsen’s conjecture on the vanishing of Galois cohomology (when X is affine) [19].

In the case of CM elliptic curves (minus a point) it is easily seen to follow also from the pseudo-nullity of a natural Iwasawa module. All of these implications are rather easy once the formalism is properly set up.

It is amusing to note that these conjectures belong to what one might call ‘the structure theory of mixed motives.’ That is to say, the usual Diophantine connection for them occurs through the theory of L-functions. It is thus rather surprising that a non-linear (and non-trivial) phenomenon like Faltings’ theorem can be linked to their validity. In the manner of physicists, it is perhaps not out of place to view these implications as positive evidence for the structure theory in question.

In relation to the ‘anabelian’ philosophy, we will explain in the last section how these implications can be viewed as a working substitute for the desired implication ‘section conjecture $\Rightarrow$ Faltings’ theorem.’

1 The De Rham unipotent Albanese map

To start out, we let L be any field of characteristic zero and S be a scheme over L . Let $f : X \rightarrow S$ be a smooth scheme over S . We denote by $\text{Un}_n(X)$ the category of unipotent vector bundles with flat connection having index of unipotency $\leq n$. That is, the objects are $(\mathcal{V}, \nabla_{\mathcal{V}})$, vector bundles $\mathcal{V}$ equipped with flat connections

$$\nabla_{\mathcal{V}} : \mathcal{V} \rightarrow \Omega_{X/S} \otimes \mathcal{V}$$

that admit a filtration

$$\mathcal{V} = \mathcal{V}_n \supset \mathcal{V}_{n-1} \supset \cdots \supset \mathcal{V}_1 \supset \mathcal{V}_0 = 0$$

by sub-bundles stabilized by the connection, such that

$$(\mathcal{V}_{i+1}/\mathcal{V}_i, \nabla) \simeq f^*(W_i, \nabla_i),$$

for some bundles with connection (W_i, ∇_i) on S . The morphisms are maps of sheaves preserving the connection. Obviously, $\text{Un}_n(X)$ is included in $\text{Un}_m(X)$ as a full subcategory if $m \geq n$, and we denote by $\text{Un}(X)$ the corresponding union. Let $b \in X(S)$ be a rational point. It determines a fiber functor $e_b : \text{Un}(X) \rightarrow \text{Vect}_S$ to the category of vector bundles on S . $\text{Un}(X)$ forms a Tannakian category, and we denote by $\langle \text{Un}_n(X) \rangle$ the Tannakian subcategory of $\text{Un}(X)$ generated by $\text{Un}_n(X)$.

Now let X be defined over L . Given any L -scheme S we follow the standard notation of X_S for the base-change of X to S . The points b and x in $X(L)$ then determine fiber functors $e_b(S), e_x(S) : \text{Un}(X_S) \rightarrow \text{Vect}_S$. We will use the notation $e_b^n(S)$, etc. to denote the restriction of the fiber functors to $\text{Un}_n(X_S)$. The notation $\langle e_b^n(S) \rangle$ will be used to denote the restriction to the category $\langle \text{Un}_n(X) \rangle$, The *De Rham fundamental group* [9] $\pi_{1,DR}(X, b)$ is the pro-unipotent pro-algebraic group over L that represents the functor on L -schemes S :

$$S \mapsto \text{Aut}^{\otimes}(e_b(S))$$

and the path space $\pi_{1,DR}(X; x, b)$ represents

$$S \mapsto \text{Isom}^{\otimes}(e_b(S), e_x(S))$$

We recall that the isomorphisms in the definition are required to respect the tensor-product structure, i.e., if $g \in \text{Isom}(e_b(S), e_x(S))$ and V, W are objects in $\text{Un}(X_S)$, then $g(v \otimes w) = g(v) \otimes g(w)$ for $v \in V_b$, $w \in W_b$. It will be convenient to consider also the L -module $\text{Hom}(e_b, e_x)$ and the L -algebra $\text{End}(e_b)$.

We will fix a base point b and denote the fundamental group by U^{DR} and the path space by $P^{DR}(x)$. All these constructions are compatible with base-change. Given three points $x, y, z \in X$, there is a ‘composition of paths’ map

$$\pi_{1,DR}(X; z, y) \times \pi_{1,DR}(X; y, x) \rightarrow \pi_{1,DR}(X; z, x)$$

that induces an isomorphism

$$\pi_{1,DR}(X; z, y) \simeq \pi_{1,DR}(X; z, x)$$

whenever one picks a point $p \in \pi_{1,DR}(X; y, x)$ (if it exists). There is an obvious compatibility when one composes three paths in different orders. In particular, $P^{DR}(x)$ naturally has the structure of a right torsor for U^{DR} . We denote by A^{DR} and $\mathcal{P}^{DR}(x)$ the coordinate rings of U^{DR} and $P^{DR}(x)$, respectively. As described in [9], section 10, the $P^{DR}(x)$ fit together to form the *canonical torsor*

$$P^{DR} \rightarrow X$$

which is a right torsor for $X \times_L U^{DR}$ and has the property that the fiber over a point $x \in X(L)$ is exactly the previous $P^{DR}(x)$. We denote by $\mathcal{P}^{DR}$ the sheaf of algebras over X corresponding to the coordinate ring of P^{DR} . In order to describe this coordinate ring, it will be convenient to describe a universal pro-unipotent pro-bundle associated to the canonical torsor.

The equivalence (loc. cit.) from the category of representations of U^{DR} to unipotent connections on X can be described as

$$V \mapsto \mathcal{V} := (P^{DR} \times V)/U^{DR}$$

Since $P^{DR}(b)$ is canonically isomorphic to U^{DR} , $\mathcal{V}_b$ is canonically isomorphic to V . Now let E be the universal enveloping algebra of $\text{Lie}U^{DR}$. Then E has the structure of a co-commutative Hopf algebra and U^{DR} is realized as the group-like elements in E . That is, if $\Delta : E \rightarrow E \hat{\otimes} E$ denotes the co-multiplication of E , then U^{DR} is canonically isomorphic to $g \in E$ such that $\Delta(g) = g \otimes g$ (of course, as we vary over points in L -algebras). Let U^{DR} act on E on the left, turning E into a pro-representation of U^{DR} : if $I \subset E$ denotes the kernel of the co-unit of E , then E is considered as the projective system of the finite-dimensional representations $E[n] := E/I^n$. Define the universal pro-unipotent pro-bundle with connection on X as

$$\mathcal{E} := (P^{DR} \times E)/U^{DR}$$

which is thus given by the projective system

$$\mathcal{E}[n] := (P^{DR} \times E[n])/U^{DR}.$$

Then $\mathcal{E}$ is characterized by the following universal property:

If $\mathcal{V}$ is a unipotent vector bundle with connection and $v \in \mathcal{V}_b$, then there exists a unique map

$$\phi_v : \mathcal{E} \rightarrow \mathcal{V}$$

such that $1 \in \mathcal{E}_b \rightarrow v \in \mathcal{V}_b$.

To see this note that $\mathcal{V}$ is associated to a representation V and that a map $\mathcal{E} \rightarrow \mathcal{V}$ is determined by a map $E = \mathcal{E}_b \rightarrow \mathcal{V}_b \rightarrow V$ of representations. But then, if $1 \mapsto v$, then $f \in E$ must map to $fv \in V$, so that the map is completely determined by the image of 1. To reformulate, there is a natural isomorphism

$$\text{Hom}(\mathcal{E}, \mathcal{V}) \simeq \mathcal{V}_b.$$

By the definition of group-like elements, the co-multiplication Δ is a map of U^{DR} representations. Therefore, there is a map of connections

$$\Delta : \mathcal{E} \rightarrow \mathcal{E} \otimes \mathcal{E}$$

which turns $\mathcal{E}$ into a sheaf of co-commutative co-algebras. This map can be also characterized as the unique map $\mathcal{E} \rightarrow \mathcal{E} \otimes \mathcal{E}$ that takes $1 \in \mathcal{E}_b$ to $1 \otimes 1 \in (cE \otimes E)_b$. We note that there is a map

$$\mathcal{E}_x \rightarrow \text{Hom}(e_b, e_x)$$

defined as follows. Suppose $\mathcal{V}$ is a unipotent vector bundle with connection, $v \in \mathcal{V}_b$ and $f \in \mathcal{E}_x$. Then $f \cdot v := (\phi_v)_x(f) \in V_x$. It is straightforward to check that this map is linear in v and functorial in $\mathcal{V}$. One also checks that all functorial homomorphisms $h : \mathcal{V}_b \rightarrow \mathcal{V}_x$ arise in this way: Given such an h , consider its value $f \in \mathcal{E}_x$ on $1 \in \mathcal{E}_b$. (There is an obvious way to evaluate h on a pro-unipotent vector bundle with connection. Here as in other places, we are being somewhat sloppy with this passage.) Now given any other $v \in \mathcal{V}$, we have $\phi_v : \mathcal{E} \rightarrow \mathcal{V}$ described above. By functoriality, we must have a commutative diagram

$$\begin{array}{ccc} \mathcal{E}_b & \xrightarrow{h} & \mathcal{E}_x \\ \phi_v \downarrow & & \downarrow \phi_v \\ \mathcal{V}_b & \xrightarrow{h} & \mathcal{V}_x \end{array}$$

Hence, we have

$$h(v) = h(\phi_v(1)) = \phi_v(h(1)) = \phi_v(f) = f \cdot v,$$

proving that the value of h is merely the action of f . Exactly the same argument with e_b^n, e_x^n in place of e_b, e_x shows that $\mathcal{E}[n]_x$ represents $\text{Hom}(e_b^n, e_x^n)$.

Now define

$$\mathcal{P} := \mathcal{E}^*,$$

which therefore is an ind-unipotent vector bundle with connection. The co-multiplication on $\mathcal{E}$ dualizes to endow $\mathcal{P}$ with the structure of a sheaf of commutative algebras over X . Since $P^{DR}(x)$ consists of the tensor compatible elements in $\text{Hom}(e_b, e_x) = \mathcal{E}_x$, we see that this corresponds to the group-like elements in $\mathcal{E}_x$ with respect to the co-multiplication, and hence, the algebra homomorphisms $\mathcal{P}(x) \rightarrow L$. Although we've carried out this argument pointwise, it applies uniformly to the whole family as follows: An argument identical to the pointwise one gives us maps

$$\mathcal{E} \rightarrow \underline{\text{Hom}}(\mathcal{V}_b \otimes \mathcal{O}_X, \mathcal{V})$$

of sheaves, that together induce an isomorphism of sheaves

$$\mathcal{E} \simeq \underline{\text{Hom}}(e_b \otimes \mathcal{O}_X, Id).$$

By definition, P^{DR} represents

$$\underline{\text{Isom}}^{\otimes}(e_b \otimes \mathcal{O}_X, Id).$$

We conclude therefore that $P^{DR} = \text{Spec}_X(\mathcal{P})$ and $\mathcal{P} = \mathcal{P}^{DR}$.

Given any pro-algebraic group G , we denote by $Z^n G$ its descending central series normalized by the indexing $Z^1 G = G$, $Z^{n+1} G = [G, Z^n G]$. Corresponding to this, we have the quotient groups, $G_n := G/Z^n G$. When the reference to the group is clear from the context, we will often omit it from the notation and write Z^n for $Z^n G$. A similar convention will apply to the various other filtrations occurring the paper. Another convenient convention we may as well mention here is that we will often omit the connection from the notation when referring to a bundle with connection. That is, $(\mathcal{V}, \nabla_{\mathcal{V}})$ will often be denoted simply by $\mathcal{V}$.

In the case of U^{DR} , we define $P_n^{DR}(x)$ to be the U_n^{DR} torsor obtained by push-out

$$P_n^{DR}(x) = (P^{DR}(x) \times U_n^{DR})/U^{DR}$$

Of course it turns out that $P_n^{DR}(x)$ represents the functor $\text{Isom}^{\otimes}(\langle e_b^n \rangle, \langle e_x^n \rangle)$. The pushout construction can be applied uniformly to the canonical torsor P^{DR} to get P_n^{DR} which is a U_n^{DR} -torsor over X .

If we define $\mathcal{P}^{DR}[n] := \mathcal{E}[n]^*$, it gives a filtration

$$\mathcal{P}^{DR}[0] = \mathcal{O}_X \subset \mathcal{P}^{DR}[1] \subset \mathcal{P}^{DR}[2] \subset \dots$$

of $\mathcal{P}^{DR}$ by finite-rank sub-bundles that we refer to as the *Eilenberg-MacLane filtration*. Observe that the multiplication

$$\mathcal{E} \otimes E \rightarrow \mathcal{E}$$

is defined by taking $f \in E$ and associating to it the unique homomorphism $\phi_f : \mathcal{E} \rightarrow \mathcal{E}$. In particular, when we compose with the projection to $\mathcal{E}[n]$, it factors through to a map

$$\mathcal{E}[n] \otimes E[n] \rightarrow \mathcal{E}[n].$$

Hence, the torsor map

$$\mathcal{P}^{DR} \rightarrow \mathcal{P}^{DR} \otimes A^{DR}$$

carries $\mathcal{P}^{DR}[n]$ to $\mathcal{P}^{DR}[n] \otimes A^{DR}[n]$. Therefore, if P^{DR} is trivialized by a point $p \in P^{DR}(S)$ in some X -scheme S , then the induced isomorphism

$$i_p : \mathcal{P}^{DR} \otimes \mathcal{O}_S \simeq A^{DR} \otimes \mathcal{O}_S$$

is compatible with the Eilenberg-MacLane filtration. We should also note that if we embed L into $\mathbb{C}$ and view the whole situation over the complex numbers, then the Eilenberg-MacLane filtration is the

one induced by the length of iterated integrals [17]. In particular, for any complex point $x \in X(\mathbb{C})$, the map obtained by base-change $\mathcal{P}^{DR}[n]_x \rightarrow \mathcal{P}_x^{DR}$ is injective. Thus, the inclusion $\mathcal{P}^{DR}[n]_x \rightarrow \mathcal{P}_x^{DR}$ is universally injective. (Proof: Given an X -scheme $f : Z \rightarrow X$, any point $z \in Z$ lies over $f(z) \in X$. So injectivity of

$$f^* \mathcal{P}^{DR}[n]_z \rightarrow f^* \mathcal{P}_z^{DR}$$

reduces to that of

$$\mathcal{P}^{DR}[n]_{f(z)} \rightarrow \mathcal{P}_{f(z)}^{DR},$$

which then can be handled through a complex embedding.) Therefore, by the constancy of dimension, we see that the trivializing map i_p just mentioned must be strictly compatible with the Eilenberg-MacLane filtration.

Now we let $L = F_v$, the completion of a number field F at a non-archimedean place v and R_v the ring of integers in F_v . Let X_v be a smooth curve over F_v . We will assume that we have a diagram

$$\begin{array}{ccccc} X_v & \hookrightarrow & \mathcal{X}_v & \hookrightarrow & \mathcal{X}'_v \\ \downarrow & & \downarrow & & \downarrow \\ \text{Spec}(F_v) & \hookrightarrow & \text{Spec}(R_v) & = & \text{Spec}(R_v) \end{array}$$

where $\mathcal{X}'_v$ is a smooth proper curve over R_v and $\mathcal{X}_v$ is the complement in $\mathcal{X}'_v$ of a smooth divisor $\mathcal{D}_v$ and X_v is the generic fiber of $\mathcal{X}_v$. We also denote by X'_v (resp. D_v) the generic fiber of $\mathcal{X}'_v$ (resp. $\mathcal{D}_v$). Letting k denote the residue field of F_v , we get varieties Y and Y' over k as the special fibers of $\mathcal{X}_v$ and $\mathcal{X}'_v$, respectively. Given points $b, x \in \mathcal{X}_v(R_v)$, we let $\bar{b}$ and $\bar{x}$ be their reduction to k -points of Y . Associated to Y and a point $c \in Y(k)$, we have the *crystalline fundamental group* $\pi_{1,cr}(Y, c)$ [6] which is a pro-unipotent pro-algebraic group over K , the field of fraction of the ring of Witt vectors W of k . The definition of this group uses the category $\text{Un}(Y)$ of over-convergent unipotent isocrystals [1] on Y which, upon base-change to F_v , can be interpreted as vector bundles $(\mathcal{V}, \nabla)$ with connections on X_v satisfying the unipotence condition of the first paragraph, a convergence condition on each residue disk $]c[$ of point $c \in Y$, and an over-convergence condition near the points of D_v . For each point $c \in Y(k)$, we get the fiber functor $e_c : \text{Un}(Y) \rightarrow \text{Vect}_K$ which, again upon base-change to F_v , associates to a pair $(\mathcal{V}, \nabla)$ the set of flat sections $\mathcal{V}(\mathcal{V})^{\nabla=0}$ on the residue disk $]c[$ ([2], p. 26). $\pi_{1,cr}(Y, c)$ then represents the group of tensor automorphisms this fiber functor. Similarly, using two points and the functor $\text{Isom}^{\otimes}(e_c, e_y)$, we get the $\pi_{1,cr}(Y, c)$ -torsor $\pi_{1,cr}(Y; y, c)$ of crystalline paths from c to y . As before, we will fix a c in the discussion and use the notation $U^{cr} = \pi_{1,cr}(Y, c)$, $P^{cr}(y) = \pi_{1,cr}(Y; y, c)$. In fact, if we let $c = \bar{b}$ and $y = \bar{x}$, then we have canonical isomorphisms ([6], prop. 2.4.1)

$$U^{DR} \simeq U^{cr} \otimes_K F_v$$

$$P^{DR}(x) \simeq P^{cr}(\bar{x}) \otimes_K F_v$$

which are compatible with the torsor structures and, more generally, composition of paths. The key point is that all algebraic unipotent bundles satisfy the (over-)convergence condition. Because $P^{cr}(y)$ is functorial, it carries a pro-algebraic automorphism

$$\phi : P^{cr}(y) \simeq P^{cr}(y)$$

induced by the $q = |k|$ -power map on $\mathcal{O}_Y$. The various Frobenius maps are also compatible with the torsor structures. The comparison isomorphism then endows P^{DR} with a map that we will again denote by ϕ . Given a point $x \in X_v(R_v)$ that lies in the residue disk $]\bar{b}[$, we have

$$P^{DR}(x) \simeq U^{cr} \otimes_K F_v$$

and it is worth chasing through the definition to see precisely what the path is on the left-hand side corresponding to the identity in U^{cr} : Given a unipotent bundle $(\mathcal{V}, \nabla)$, we construct an isomorphism $\mathcal{V}_b \simeq \mathcal{V}_x$ by finding the unique flat section s over $]\bar{b}[$ with initial value at b equal to v_b . Then the

image of v_b under the isomorphism (i.e., the path) is $s(x)$, the value of this flat section at x . Since ϕ also respects the subcategories $\text{Un}_n(Y)$ consisting of the overconvergent isocrystals with index of unipotency $\leq n$, it preserves the Eilenberg-MacLane filtration on the $\mathcal{P}^{DR}(x)$.

According to [29], theorem E, $\mathcal{P}^{DR}$ possesses a Hodge filtration. This is a filtration

$$\mathcal{P}^{DR} = F^0 \mathcal{P}^{DR} \supset \dots \supset F^n \mathcal{P}^{DR} \supset F^{n+1} \mathcal{P}^{DR} \supset \dots$$

by sub- $\mathcal{O}_X$ -modules that is compatible with the multiplicative structure. In particular, the F^i are ideals. This induces a filtration on the fibers $\mathcal{P}^{DR}(x)$ and in particular on the coordinate ring A^{DR} of U^{DR} . The Hodge filtration is compatible with the torsor structure in that if the tensor product $\mathcal{P}^{DR} \otimes_L A^{DR}$ is endowed with the tensor product filtration

$$F^n(\mathcal{P}^{DR} \otimes_L A^{DR}) = \sum_{i+j=n} (F^i \mathcal{P}^{DR}) \otimes_L (F^j A^{DR})$$

then the torsor map

$$\mathcal{P}^{DR} \rightarrow \mathcal{P}^{DR} \otimes_L A^{DR}$$

is compatible with the filtration. As in the case of the Eilenberg-MacLane filtration, if we utilize their descriptions over $\mathbb{C}$ [18], we see immediately that the inclusions $F^i \mathcal{P}^{DR} \hookrightarrow \mathcal{P}^{DR}$ are also universally injective, and that if we restrict the Hodge filtration to the terms in the Eilenberg-MacLane filtration, then the rank of $F^i \mathcal{P}^{DR}[n]$ is equal to the dimension of $F^i A^{DR}[n]$. Since the F^i are ideals, there are corresponding sub- X -schemes

$$F^i P^{DR},$$

where $F^{-i+1} \mathcal{P}^{DR}$ is the defining ideal for $F^i P^{DR}$. In fact, $F^0 P^{DR}$ is a $F^0 U^{DR}$ -torsor. To see the action, note that compatibility implies that the torsor map takes $F^1 \mathcal{P}^{DR}$, the defining ideal for $F^0 P^{DR}$, to

$$F^1 \mathcal{P}^{DR} \otimes A^{DR} + \mathcal{P}^{DR} \otimes F^1 A^{DR},$$

the latter being exactly the defining ideal for $F^0 P^{DR} \times F^0 U^{DR}$ inside $P^{DR} \times U^{DR}$. Thus, we get the action map

$$F^0 P^{DR} \rightarrow F^0 P^{DR} \times F^0 U^{DR}.$$

On the other hand, if we have a point $p \in F^0 P^{DR}(S)$ in some X -scheme S . Then the corresponding isomorphism

$$i_p : \mathcal{P}^{DR} \otimes \mathcal{O}_S \simeq A_{DR} \otimes \mathcal{O}_S$$

is compatible with the Hodge filtration and hence, strictly compatible because of the universal injectivity mentioned above and the equality of ranks (of course when restricted to the finite-rank terms of the Eilenberg-MacLane filtration). Therefore, it induces an isomorphism

$$i_p : F^0 U^{DR} \simeq F^0 P^{DR},$$

proving the torsor property. (The remaining compatibilities are obvious from the corresponding properties for P^{DR} .) Regarding these trivializations, we note that if a point $p \in P^{DR}(S)$ induces an isomorphism as above compatible with the Hodge filtration, then since p is recovered as $ev_0 \circ i_p$, where $ev_0 : A_{DR} \otimes \mathcal{O}_S \rightarrow \mathcal{O}_S$ is the origin of U_S^{DR} , we see that $p \in F^0 P^{DR}$.

We will need some abstract definitions corresponding to the situation described. Given an F_v -scheme Z , by a torsor over Z for U^{DR} , we mean a right U_Z^{DR} torsor $T = \text{Spec}(T)$ over Z endowed with an ‘Eilenberg-MacLane’ filtration:

$$\mathcal{T}[0] \subset \mathcal{T}[1] \subset \dots$$

and a ‘Hodge’ filtration

$$\mathcal{T} = F^0 \mathcal{T} \supset F^1 \mathcal{T} \supset \dots$$

of the coordinate ring. The Eilenberg-MacLane filtration should consist of locally-free $\mathcal{O}_Z$ -modules of finite rank equal to $\dim_{F_v} A^{DR}[n]$ with the property that $\mathcal{T}[n] \hookrightarrow \mathcal{T}$ is universally injective. Furthermore, each $F^i \mathcal{T}[n] \hookrightarrow \mathcal{T}[n]$ must be universally injective and have rank equal to $\dim F^i A^{DR}[n]$. Both filtrations are required to be compatible with the torsor structure in slightly different senses, namely, that $\mathcal{T}[n]$ must be carried to $\mathcal{T}[n] \otimes A^{DR}[n]$ while $F^m \mathcal{T}[n]$ is to be taken to $\Sigma_{i+j=m} F^i \mathcal{T} \otimes F^j A^{DR}$. The torsor T must also carry a ‘Frobenius’ automorphism of Z -schemes $\phi : T \rightarrow T$ (we will denote all of them by the same letter) which is required to be compatible with the torsor structure in the sense that $\phi(t)\phi(u) = \phi(tu)$ for points t of T and u of U . We require that ϕ preserves the Eilenberg-MacLane filtration, although not necessarily the Hodge filtration. We will call the torsor T *admissible*, if it is separately trivializable for the two Frobenius structure and the Hodge filtration. That is, we are requiring that there is a point $p^{cr} \in T(Z)$ which is invariant under the Frobenius, and also that there is a point $p^{DR} \in F^0 T(Z)$.

The following lemma follows from the argument of [2], cor. 3.2.

Lemma 1 *Let T any torsor for U^{DR} over an affine scheme Z . Then T is uniquely trivializable with respect to the ϕ -structure. That is, there is a unique point $p_T^{cr} \in T(Z)$ which is invariant under ϕ .*

The key fact is that the map $U^{DR} \rightarrow U^{DR}$, $x \mapsto \phi(x^{-1})x$ is surjective. Obviously, the canonical path is the identity for the trivial torsor U_Z^{DR} .

Combining this lemma with the earlier remarks and the unipotence of $F^0 U^{DR}$, we get that for any affine X -scheme Z , P_Z^{DR} is an admissible torsor.

We say two admissible torsors are isomorphic if there is a torsor isomorphism between them that is simultaneously compatible with the Frobenius and Hodge filtration.

Let T be an admissible U^{DR} torsor over an F_v -algebra L . Choose a trivialization $p_T^H \in F^0 T(L)$. There then exists a unique element $u_T \in U^{DR}$ such that $p_T^{cr} u_T = p_T^H$. Clearly, a different choice of p^H will change the result only by right multiplication with an element of $F^0 U^{DR}$. Thus, we get an element $[u_T] \in U^{DR}/F^0 U^{DR}$ which is independent of the choice of p_T^H .

Proposition 1 *The map*

$$T \mapsto [u_T]$$

defines a natural bijection from the isomorphism classes of admissible torsors to $U^{DR}/F^0 U^{DR}$. That is to say, the scheme $U^{DR}/F^0 U^{DR}$ represents the functor that assigns to each F_v algebra L the isomorphism classes of admissible torsors on L .

Proof. The map defined is clearly functorial, so we need only check bijectivity on points. Suppose $[u_T] = [u_S]$. Then there exists a $u^0 \in F^0 U^{DR}$ such that $u_S = u_T u^0$. The elements p_T^{cr} and p_S^{cr} already determine ϕ -compatible isomorphisms

$$f : S \simeq U^{DR} \simeq T; \quad p_S^{cr} u \mapsto u \mapsto p_T^{cr} u$$

It suffices to check that this isomorphism is compatible with the Hodge filtration. But we already know that the map

$$h : p_S^H g \mapsto g \mapsto p_T^H g$$

is compatible with the Hodge filtration, and writing the ϕ -compatible map f with respect to the p^H ’s, we get

$$p_S^H g = p_S^{cr} u_S g \mapsto p_T^{cr} u_S g = p^H u_T^{-1} u_S g = p^H u^0 g.$$

That is, it corresponds merely to a different choice $p^H u^0$ of the trivialization. So we need to check that $p^H u^0 \in F^0 T$, i.e., that it is a Hodge trivialization. Its values on the coordinate ring of T is expressed through the composition

$$\mathcal{T} \rightarrow \mathcal{T} \otimes A^{DR} \xrightarrow{p^H \times u^0} L$$

But then, since the action takes $F^1\mathcal{T}$ to

$$F^1\mathcal{T} \otimes A^{DR} + \mathcal{T} \otimes F^1A^{DR},$$

$F^1\mathcal{T}$ is clearly killed by this evaluation, i.e., $p^Hu^0 \in F^0T$.

Thus the map of functors described is injective. To see that it is surjective, note that for any Z , we have

$$U^{DR}/F^0U^{DR}(Z) = U^{DR}(Z)/F^0U^{DR}(Z).$$

Now, given $g \in U^{DR}$, we consider U^{DR} with the same Hodge filtration, but with the automorphism twisted to $\phi^g(h) = g^{-1}\phi(gh)$. This automorphism is compatible with right multiplication, that is, the right torsor structure on U^{DR} :

$$\phi^g(hk) = g^{-1}\phi(ghk) = g^{-1}\phi(gh)\phi(k) = \phi^g(h)\phi(k)$$

Thus, we end up with an admissible torsor T_g . The ϕ^g -fixed element here is clearly g^{-1} , and thus, $u_{T_g} = g$, as desired. $\square$

Already we have the language necessary to define the De Rham unipotent Albanese map:

$$j^{DR}\mathcal{X}_v(R_v) \rightarrow U^{DR}/F^0U^{DR}$$

by

$$j^{DR}(x) := [P^{DR}(x)].$$

The finite-level versions j_n^{DR} are defined by composing with the natural projections

$$U^{DR}/F^0U^{DR} \rightarrow U_n^{DR}/F^0U_n^{DR}.$$

There is a parallel discussion of admissible torsors for U_n^{DR} out of which one can extract the interpretation of

$$j_n^{DR} : \mathcal{X}_v(R_v) \rightarrow U_n^{DR}/F^0U_n^{DR}$$

as

$$x \mapsto [P_n^{DR}(x)]$$

In order to describe the map, it will be convenient to have a rather explicit construction of U^{DR} and P^{DR} . We will carry this out assuming X_v is affine, which is the only case we will need. The construction depends on the choice $\alpha_1, \alpha_2, \dots, \alpha_m$ of global algebraic differential forms representing a basis of $H_{DR}^1(X_v)$. So here, we have $m = 2g + s - 1$, where s is the order of $X'_v \setminus X_v$. Corresponding to this choice, there is a free non-commutative algebra

$$F_v < A_1, \dots, A_m >$$

generated by the symbols $A_1, A_2, \dots, A_m$. Thus, it is the group algebra of the free group on m generators. Let I be its augmentation ideal. The algebra $F_v < A_1, \dots, A_m >$ has a natural comultiplication map Δ with values $\Delta(A_i) = A_i \otimes 1 + 1 \otimes A_i$. Now we consider the completion

$$E := \varprojlim F_v < A_1, \dots, A_m > / I^n$$

Δ extends naturally to a comultiplication $E \rightarrow E \hat{\otimes} E$. We can also consider the quotients

$$E[n] := E / I^{n+1}A = F_v < A_1, \dots, A_m > / I^{n+1}$$

that carry the induced comultiplication. Thus, E and E_n have the structure of non-commutative, co-commutative Hopf algebras.

Now let $\mathcal{E}$ be the pro-unipotent pro-vector bundle $E \otimes \mathcal{O}_{X_v}$ with the connection ∇ determined by

$$\nabla f = -\sum_i A_i f \alpha_i$$

for constant sections $f \in E$. We also have the finite-level quotients $\mathcal{E}[n] := E[n] \otimes \mathcal{O}_{X_v}$. This construction ends up as the ‘universal’ pro-unipotent pro-bundle with connection in the sense of the beginning paragraphs, justifying the conflation of notation. To see this, we need a lemma.

Lemma 2 *Let (V, ∇) be a unipotent bundle with flat connection on X_v of rank r . Then there exist strictly upper-triangular matrices N_i such that*

$$(V, \nabla) \simeq (\mathcal{O}_X^r, d + \sum_i \alpha_i N_i)$$

Proof of lemma. Since X_v is affine and (V, ∇) is unipotent, by choosing vector bundle splittings of the filtration, there exists a trivialization $V \simeq \mathcal{O}_X^r$ such that ∇ takes the form $d + \omega$ for some strictly upper-triangular $n \times n$ matrix of 1-forms ω . It will be convenient to write $\omega = \sum \omega_{ij} E_{ij}$ where E_{ij} is the elementary $n \times n$ matrix with a 1 in the (i, j) -entry and zero elsewhere, and $\omega_{ij} = 0$ unless $j > i$. Recall that a gauge transformation G will change the connection matrix by

$$\omega \mapsto G^{-1} \omega G + G^{-1} dG$$

For a gauge transformation of the form $G = I - aE_{ij}$ ($j \neq i$), we have $G^{-1} = I - aE_{ij}$ while $G^{-1}dG = daE_{ij}$. We wish to perform a series of gauge transformations so that each entry ω_{ij} is replaced by a linear combination of the α_i 's and β_j 's. Now, any single one-form can be written as such a combination after adding an exact form. We will show how to change ω_{ij} by induction on $j - i$. Assume we are done for $j - i < c$. So let i, j satisfy $j - i = c$. First find a such $\omega_{ij} + da$ is a linear combination of the α_i and β_j . Now consider the gauge transformation $G = I - aE_{ij}$. We compute

$$G^{-1} \omega = (I + aE_{ij})(\sum \omega_{kl} E_{kl}) = \omega + \sum_l \omega_{jl} a E_{il} =: \omega'$$

Note the only non-zero l occurring in the sum are strictly bigger than j . Thus, $\omega'_{\mu\nu} \neq \omega_{\mu\nu}$ only occurs for $\mu = i$ and $\nu > j$, that is, for $\nu - \mu > j - i$. On the other side,

$$\omega' G = (\sum \omega'_{kl} E_{kl})(I - aE_{ij}) = \omega' - \sum \omega'_{ki} a E_{kj}$$

and all the k appearing in the sum are $< i$. Therefore, we see that all the entries ω'_{kl} of $\omega^G := G^{-1} \omega G$ are equal to ω_{kl} for $l - k \leq c$. On the other hand, $dGG^{-1} = daE_{ij}$. Performing such a gauge transformation for each i, j such that $j - i = c$ clearly achieves what we want. $\square$.

Now we can formulate the universal property of $\mathcal{E}$:

Lemma 3 *Given any object (V, ∇) in $Un(X_v)$ together with a element $v \in V_b$ (the fiber at b), there exists a unique morphism $\phi : \mathcal{E} \rightarrow V$ in $Un(X_v)$ such that $1 \in \mathcal{E}_b \mapsto v$. That is, $\mathcal{E}$ is the same as the bundle with the same notation from the beginning of the section.*

Proof. First we will show uniqueness. For this, we can choose an embedding of F_v into the complex numbers and assume that everything is defined over $\mathbb{C}$ (and for convenience, we will omit the base-change to $\mathbb{C}$ from the notation). By uniqueness of solutions to differential equations with given initial condition, we see that any map of bundles $\phi : \mathcal{E} \rightarrow V$ is completely determined by its value at b . (Actually, for this, we needn't go to $\mathbb{C}$. It suffices to use uniqueness of formal solutions.) So we need to check that the value of the map is determined at b by the given conditions. Let π be the topological fundamental group of $X_v(\mathbb{C})$ and let $\mathbb{C}[\pi]$ be the complex group algebra of π . The holonomy transformations give an action of $\mathbb{C}[\pi]$ on the fibers $\mathcal{E}_b$ and V_b . Since ϕ respects the connections, ϕ_b is equivariant for this action. But over $\mathbb{C}$, one can compute the holonomy using iterated integrals. Given a word $w = A_{i_1} A_{i_2} \cdots A_{i_n}$ in the A_i and a piece-wise smooth path γ , denote by $\int_\gamma \alpha_w$, the iterated integral

$$\int_\gamma \alpha_{i_1} \alpha_{i_2} \cdots \alpha_{i_n}.$$

We are using the normalization whereby if $\gamma^*(\omega_i) = f_i(t)dt$, then

$$\int_\gamma \omega_1 \omega_2 \cdots \omega_n = \int_0^1 f_1(t_1) \int_0^{t_1} f_2(t_2) \int_0^{t_2} f_3(t_3) \int \cdots \int_0^{t_{n-1}} f_n(t_n) dt_n dt_{n-1} \cdots dt_1.$$

On a contractible open set U , if we pick a base point c and define the function $G_c(z)$ with values in E on U by

$$G_c(z) = \Sigma_w \int_c^z \alpha_w[w],$$

where the iterated integral occurs along any path from c to z , then $dG_c = \Sigma_i \alpha_i A_i G_c$ so that $G_c(z)$ can be viewed as a flat section of $\mathcal{E}$. In particular, for the holonomy around a loop γ based at b ,

$$\gamma \cdot f = \Sigma_w \int_\gamma \alpha_w[w] f$$

for $f \in E = \mathcal{E}_b$. Thus, we must have

$$\phi(\Sigma_w \int_\gamma \alpha_w[w] 1) = \gamma \phi(1) = \gamma v$$

Now consider the map $\mathbb{C}[\pi] \rightarrow E$ given by the formula

$$\gamma \mapsto G_\gamma := \Sigma_w \int_\gamma \alpha_w[w]$$

(We will use the natural notation $\int_\gamma \alpha_w := \Sigma_i c_i \int_{\gamma_i} \alpha_w$ if $\gamma = \Sigma_i c_i \gamma_i$, $c_i \in \mathbb{C}$, $\gamma_i \in \pi$.) To conclude the proof of uniqueness, we just need to see that the composition to each of the finite-dimensional quotients E_n is surjective. But if not, we would have a non-trivial linear relation between the $\int_\cdot \alpha_w$ regarded as functions on $\mathbb{C}[\pi]$. We show by induction on the length of w that they are in fact linearly independent. For this, let $a_1, \dots, a_m$ be elements of $\mathbb{C}[\pi]$ whose classes in homology form a dual basis to the α_i . Assume

$$\Sigma_{|w| \leq n} c_w \int_\gamma \alpha_w = 0$$

for all $\gamma \in \pi$. Write this as

$$\Sigma_{|w| \leq n-1} (\Sigma_i c_{A_i w} \int_\gamma \alpha_{A_i w}) = 0$$

and apply it to $a_j \gamma$. Then we have

$$\Sigma_{|w| \leq n-1} (\Sigma_i c_{A_i w} \int_{a_j \gamma} \alpha_i \alpha_w) = 0$$

The co-product formula for iterated integrals [5] can then be applied to give

$$\Sigma_{|w| \leq n-1} [\Sigma_i c_{A_i w} (\deg(a_j) \int_\gamma \alpha_i \alpha_w + \int_{a_j} \alpha_i \int_\gamma \alpha_w + \dots)] = 0,$$

where $\deg(\Sigma c_\gamma [\gamma]) = \Sigma c_\gamma$ on elements of $\mathbb{C}[\pi]$. From this, we subtract $(\deg(a_j))$ times the original relation to get

$$\Sigma_{|w|=n-1} c_{A_j w} \int_\gamma \alpha_w + \Sigma_{|w| \leq n-2} b_w \int_\gamma \alpha_w = 0$$

for some constants b_w . Since this holds for all γ , by induction, we have $c_{A_j w} = 0$ for $|w| = n-1$, and hence $c_w = 0$ for $|w| = n$ when the argument is applied to each A_j . Then again by induction, all the c_w must vanish.

The existence part is easy. We may assume that (V, ∇) is of the form $(\mathcal{O}_{X_v}^r, d - \Sigma_i \alpha_i N_i)$ as in the previous lemma, except we have changed N_i to $-N_i$. Now given the vector $v \in V_b$, define the map $\mathcal{E} \rightarrow V$ that sends a section $\Sigma f_w[w]$ of $\mathcal{E}$ to $\Sigma_w f_w N_w v$ where the notation is that if a word has the form $w = A_{i_1} A_{i_2} \dots A_{i_n}$, then N_w is the matrix $N_{i_1} N_{i_2} \dots N_{i_n}$. This map is obviously compatible with the

connection and takes the constant section 1 to the constant section v (in particular, at the point b).
 $\square$

As mentioned already, another way of phrasing the lemma is to say that there is a canonical isomorphism:

$$\mathrm{Hom}(\mathcal{E}, V) \simeq V_b$$

In particular, $\mathrm{Hom}(\mathcal{E}, \mathcal{E}) \simeq \mathcal{E}_b = E$ and it is easily checked that this isomorphism is compatible with the algebra structure.

Recall the evaluation fiber functor $e_b : \mathrm{Un}(X_v) \rightarrow \mathrm{Vect}_{F_v}$. Since $\mathcal{E}$ is the universal pro-unipotent vector bundle with connection, $E = \mathcal{E}_b$ represents $\mathrm{End}(e_b)$ and $E = \mathcal{E}_x$ represents $\mathrm{Hom}(e_b, e_x)$. It is equally easy to see that $E[n]$ represents $\mathrm{End}(e_b^n)$ and $\mathrm{Hom}(e_b^n, e_x^n)$.

If we examine the co-multiplication $\Delta : \mathcal{E} \rightarrow \mathcal{E} \otimes \mathcal{E}$ associated to the $\mathcal{E}$ as a universal object, that is, the unique map that sends 1 to $1 \otimes 1$, by the definition of the connection on $\mathcal{E}$, it takes A_i to $A_i \otimes 1 + 1 \otimes A_i$. That is, the co-multiplication is compatible with the one defined formally on $E = \mathcal{E}_b$. So we also have that $\mathrm{Aut}^\otimes(e_b)$ and $\mathrm{Isom}^\otimes(e_b, e_x)$ are represented by the group-like elements in $\mathcal{E}_b = E = \mathcal{E}_x$ and that $\mathcal{E}^*$, the dual bundle of $\mathcal{E}$, with its structure of a commutative algebra is nothing but $\mathcal{P}^{DR}$.

Now $\mathcal{P}^{DR}$ carries a Hodge filtration by sub-bundles, but its explicit description will be unnecessary for our purposes. All we will need is that $F^0 P$ is a torsor for the group $F^0 U^{DR}$, and hence, can be trivialized over X_v in the case it is affine.

Proof of theorem 1.

First note that if X_v happens to be compact, we can delete an R_v point x_0 away from the residue disk of b to get $\mathcal{X}_v^0 = \mathcal{X}_v \setminus x_0$, still equipped with an R_v point b . Since restriction embeds the unipotent bundles with flat connection on X_v as a full subcategory of the unipotent bundles on X_v^0 (see, for example, [2], cor. 2.15), we get a surjection $\pi_{1, DR}(X_v^0, b) \rightarrow \pi_{1, DR}(X_v, b)$. This map is compatible with the Frobenius because the comparison isomorphism with π_1^{cr} is functorial [6]. Functoriality of the Hodge filtration can be deduced either from an examination of Wojkowiak's construction or the usual functoriality over $\mathbb{C}$ [17]. Of course, an identical argument can be applied to the path spaces, giving us a commutative diagram

$$\begin{array}{ccc} \mathcal{X}^0(R_v) & \rightarrow & \pi_{1, DR}(X_v^0, b)/F^0 \\ \downarrow & & \downarrow \\ \mathcal{X}(R_v) & \rightarrow & \pi_{1, DR}(X_v, b)/F^0 \end{array}$$

where the right vertical map is surjective. Therefore, it suffices to prove the theorem for an affine curve.

In fact, we will show that the image of the residue disk $]\bar{b}[$ is already Zariski dense. Now, as described above, there exists a canonical U^{DR} torsor P^{DR} on X_v with the property that the fiber over x is exactly $P^{DR}(x)$ which is, in fact, admissible. This is because of the unique Frobenius invariant element and the fact that $F^0 P^{DR}$ is a $F^0 U^{DR}$ -torsor, and hence, has a section over an affine scheme. Choose such a trivialization $p^H \in F^0 P^{DR}(X_v)$. It is important to note here that $F^0 P^{DR}$ becomes thereby *algebraically* trivialized. We can compare this with any other algebraic trivialization $g \in P^{DR}(X_v)$. That is, there exists an algebraic map $\gamma : X_v \rightarrow U^{DR}$ such that $g\gamma = p^H$. For any $x \in X_v$, we have a point $u(x) \in U^{DR}$ such that $p^{cr}(x)u(x) = g(x)$ so that $p^{cr}(x)u(x)\gamma(x) = p^H(x)$. That is, $j^{DR}(x) = [u(x)\gamma(x)]$. Therefore, it suffice to show that the image of $x \mapsto u(x)\gamma(x)$ is Zariski dense in $U^{DR}(X)$. As previously discussed, given (V, ∇) , on $]\bar{b}[$, the element $p^{cr}(x)$ is obtained from the diagram

$$\begin{array}{ccc} & V(]\bar{b}[)^{\nabla=0} & \\ \simeq \swarrow & & \searrow \simeq \\ V_b & & V_x \end{array}$$

as the inverse of the left arrow followed by the right arrow. We proceed now to describe the various objects in local coordinates to obtain the desired result.

Take g to be the trivialization of P^{DR} determined by the previously described trivialization of $\mathcal{E}$. That is, since we have identified $\text{Isom}^\otimes(e_b, e_x) = E$, the element in question is $1 \in E$. For $x \in]\bar{b}[$ $u(x)$ is then identified with the value of the function $u :]\bar{b}[\rightarrow E$ that is the unique horizontal section of $\mathcal{E}$ convergent on $]\bar{b}[$ such that $u(b) = 1$. Write $u = \sum_w u_w[w]$, where w runs over the words in A_i . We will show:

Lemma 4 *The u_w are linearly independent over the algebraic functions on X_v .*

It is easy to see that this lemma implies the theorem:

We have

$$j^{DR}(x) = \sum_w j_w(x)[w] = u(x)\gamma(x) = \sum[\sigma_{w'w''=w} u_{w'}(x)\gamma_{w''}(x)][w],$$

where $\gamma = \sum_w \gamma_w[w]$. If the image of j_n^{DR} were in a Zariski closed subspace, we would have a linear relation among the j_w , by the description we have given of the coordinate ring of U^{DR} . Let us see by induction on the length of w that the j_w are linearly independent. Assume we have a relation

$$\sum_{|w| \leq n} c_w j_w = 0$$

Then

$$\sum_{|w| \leq n} c_w [\sum_{w'w''=w} u_{w'}\gamma_{w''}] = 0.$$

This would give us a relation of the form

$$\sum_{|w| \leq n} a_w u_w = 0$$

where the a_w are algebraic functions. Hence, all $a_w = 0$. But for $|w| = n$, we have $a_w = c_w \gamma_0$ and $\gamma_0 = 1$ because γ is group-like. Therefore, we must have $c_w = 0$ for all $|w| = n$. The result follows by induction.

So we need only give the

Proof of lemma. The proof here is entirely similar to the uniqueness part of Lemma 4. Even though the u_w are convergent on $]\bar{b}[$ we can regard them as formal power series and show independence over the algebraic functions in there. For this, we choose an embedding $F_v \hookrightarrow \mathbb{C}$ into the complex numbers, so that we are in the situation of a connection on a curve over $\mathbb{C}$. Uniqueness of formal solutions show that the u_w are just the power series expansions of analytic solutions near b . But the analytic solutions can be obtained as iterated integrals: in a contractible neighborhood of b , choose a path c from b to x . Then for $w = A_{i_1} A_{i_2} \cdots A_{i_n}$, we have

$$u_w = \int_c \alpha_{i_1} \alpha_{i_2} \cdots \alpha_{i_n}$$

We see therefore, that $u(x)$ can also be continued as a multi-valued function to all of X_v (which, recall, we are now regarding as a complex curve). Regard the u_w as functions on the universal covering $\tilde{X}_v$ after choosing a point $b' \in \tilde{X}_v$ lying over b . Let the topological fundamental group π of X_v act this time on the space of analytic functions on $\tilde{X}_v$ by $\gamma : f(z) \mapsto f^\gamma(z) = f(z\gamma)$. This extends to an action of the complex group algebra $\mathbb{C}[\pi]$. To compute $u_w^\gamma(z)$ for an element $\gamma \in \Gamma$, we proceed as follows: Choose a path c from $b'\gamma$ to z . Now choose a lifting $\tilde{\gamma}$ of γ to a path in $\tilde{X}_v$ from b' to $b'\gamma$. Then

$$u_w(z\gamma) = \int_{c\tilde{\gamma}} \alpha_{i_1} \cdots \alpha_{i_n} = u_w(z) + \int_c \alpha_{i_1} \cdots \alpha_{i_{n-1}} \int_{\tilde{\gamma}} \alpha_{i_n} + \int_c \alpha_{i_1} \cdots \alpha_{i_{n-2}} \int_{\tilde{\gamma}} \alpha_{i_{n-1}} \alpha_{i_{n-2}} + \cdots$$

As before, let $a_i \in \mathbb{C}[\pi]$ be elements whose image in $H_1(X_v, \mathbb{C})$ form a basis dual to the α_i . Then we see that

$$u_{wA_i}^{a_i} = u_{wA_i} + u_w + \sum_{|w'| < |w|} c_{w'} u_{w'}$$

for constants $c_{w'}$.

Now we prove the desired independence by induction on $|w|$. So assume we are done for $|w| < n$, and suppose we have a linear relation

$$\Sigma_{|w| \leq n} h_w u_w = 0$$

where the h_w are algebraic. In particular, they are single-valued on X_v , and so are acted on trivially by π . We write this relation as

$$\Sigma_{|w|=n} h_w u_w + \Sigma_{|w|=n-1} h_w u_w + \Sigma_{|w| \leq n-2} h_w u_w = 0$$

or

$$\Sigma_i [\Sigma_{|w|=n-1} h_{wA_i} u_{wA_i} + \Sigma_{|w|=n-2} h_{wA_i} u_{wA_i}] + \Sigma_{|w| \leq n-2} h_w u_w = 0$$

When we apply $a_j - I$ to this relation, we get

$$\Sigma_{|w|=n-1} h_{wA_j} u_w + \Sigma_{|w| \leq n-2} b_w u_w = 0$$

for some algebraic functions b_w . By induction, we get $h_{wA_j} = 0$ for all w . Since this works for any j , we are done. $\square$

2 Étale Realizations

This time we start with a smooth scheme X over a field L of characteristic zero and put $\bar{X} = X \otimes_L \bar{L}$. We have on $\bar{X}$ the category $\text{Un}_p^{et}(\bar{X})$ of $\mathbb{Q}_p$ -unipotent lisse sheaves. Choosing a point $b \in \bar{X}$ then determines a fiber functor $e_b : \text{Un}_p^{et}(\bar{X}) \rightarrow \text{Vect}_{\mathbb{Q}_p}$ by taking stalks. We define the pro-unipotent p -adic étale fundamental group (see, e.g., [28]) and path space by

$$\pi_{1,\text{ét}}(\bar{X}, b) = \text{Aut}^\otimes(e_b)$$

and

$$\pi_{1,\text{ét}}(\bar{X}; x, b) = \text{Isom}^\otimes(e_b, e_x),$$

where, as usual, we have represented an obvious functor on $\mathbb{Q}_p$ algebras. When both b and x are in $X(L)$, we get an action of $G = \text{Gal}(\bar{L}/L)$. In particular, $P^{et}(x) = \pi_{1,\text{ét}}(\bar{X}; x, b) = \text{Spec}(\mathcal{P}^{et})$ is a right $U^{et} = \pi_{1,\text{ét}}(\bar{X}, b) = \text{Spec}(A^{et})$ torsor in a manner compatible with the Galois action. Here as well, we can consider the subcategory $\text{Un}_{p,n}^{et}(\bar{X})$ of unipotent $\mathbb{Q}_p$ local systems V having index of unipotency $\leq n$ in that V is endowed with a filtration

$$V = V_n \supset V_{n-1} \supset \cdots \supset V_1 \supset V_0 = 0$$

such that each successively quotient is isomorphic to a trivial $\mathbb{Q}_p$ -sheaf. The corresponding restrictions of the fiber functors will again be denoted e_b^n , etc. We denote by $\langle \text{Un}_{p,n}^{et}(\bar{X}) \rangle$ the Tannakian subcategory of $\text{Un}_p^{et}(\bar{X})$ generated by $\text{Un}_{p,n}^{et}(\bar{X})$. Using the restrictions $\langle e_b^n \rangle := e_b|_{\langle \text{Un}_{p,n}^{et}(\bar{X}) \rangle}$ and $\langle e_x^n \rangle := e_x|_{\langle \text{Un}_{p,n}^{et}(\bar{X}) \rangle}$ of the fiber functors, we then get U_n^{et} and $P_n^{et}(x)$. In the above, we can everywhere consider homomorphisms of functors rather than tensor isomorphisms, like $\text{Hom}(e_b, e_x)$, which of course have the structure of $\mathbb{Q}_p$ vector spaces.

It might be useful to recall at this point some rudimentary points about the structure of these groups and torsors, although it is not necessary to be as explicit as in the De Rham case.

Let E^{et} be the universal enveloping algebra of Lie U^{et} . Then E^{et} again has the structure of a co-commutative Hopf algebra, and we define $E^{et}[n] := E^{et}/I^n$, where $I \subset E$ is the kernel of the counit $E^{et} \rightarrow \mathbb{Q}_p$. As in the De Rham situation, we have the universal pro-unipotent lisse $\mathbb{Q}_p$ -sheaf $\mathcal{E}^{et}$ that can be constructed, for example, by twisting the canonical profinite torsor $\hat{T} := P^\wedge|X \times b$, in the notation of [9], section 10.17, with representation of the profinite $\hat{\pi}_1(\bar{X}, b)$ obtained by composing the natural map

$$\hat{\pi}_1(\bar{X}, b) \rightarrow U^{et}$$

with the left multiplication of U^{et} on E . By an argument identical to the De Rham setting, we deduce the property that $\mathcal{E}_x^{et}$ represents $\text{Hom}(e_b, e_x)$ and the finite-rank quotients $\mathcal{E}^{et}[n]$ corresponding to E/I^n represent in their turn $\text{Hom}(e_b^n, e_x^n)$. Once again the universal property gives us a comultiplication

$$\Delta : \mathcal{E}^{et} \rightarrow \mathcal{E}^{et} \otimes \mathcal{E}^{et}$$

(here and henceforward, a tensor product without an explicit subscript will refer to tensoring over $\mathbb{Q}_p$) so that if we define $\mathcal{P}^{et} = \text{Hom}(\mathcal{E}^{et}, \mathbb{Q}_p)$, then $\mathcal{P}^{et}$ is an ind- $\mathbb{Q}_p$ lisse sheaf with the property that $\mathcal{P}_x^{et}$ is the coordinate ring of $P^{et}(x)$. There is also a parallel (to the De Rham setting) discussion corresponding to $\mathcal{P}^{et}[n]$, $\mathcal{P}_n^{et}$, and P_n^{et} , which we will assume without further comment. We will refer again to the filtration by the $\mathcal{P}^{et}[n]$ as the Eilenberg-MacLane filtration. When $x \in X(L)$, since the categories $\text{Un}_{p,n}^{et}(\bar{X})$ are invariant under the Galois action, so is $\mathcal{P}^{et}[n](x)$. Therefore, $\mathcal{P}^{et}$ is a direct limit of the finite-dimensional Galois modules $\mathcal{P}^{et}[n](x)$, allowing much of the theory of finite-dimensional representations, for example Fontaine's theory of the Dieudonné functor [11], to apply.

In the subsequent discussion, fix a quotient group G through which the action of $\text{Gal}(\bar{L}/L)$ factors.

We recall some notions from [22]. Given a $\mathbb{Q}_p$ -algebra R , we give it the inductive limit topology obtained from the natural topology on finite-dimensional $\mathbb{Q}_p$ -subspaces. Then by a torsor for U^{et} over R , we mean a U_R^{et} torsor $T = \text{Spec}(\mathcal{T})$ in the usual sense, except that we require that

- (1) $\mathcal{T} = \varinjlim_n \mathcal{T}[n]$ for some locally free sub- R -modules $\mathcal{T}[n]$ that are local direct summands and of finite rank equal to $\dim A^{et}[n]$, as n runs through the non-negative integers.
- (2) $\mathcal{T}$ is equipped with a continuous R -linear action of G that stabilizes the $\mathcal{T}[n]$.
- (3) The torsor structure

$$\mathcal{T} \rightarrow \mathcal{T} \otimes_R (A^{et} \otimes R)$$

is compatible with the G -action and takes $\mathcal{T}[n]$ to $\mathcal{T}[n] \otimes_R (A^{et}[n] \otimes R)$.

When L is the completion F_v of a number field F as in the introduction and $G = G_v = \text{Gal}(F_v/F)$, we get a classifying pro-variety $H^1(G_v, U^{et})$ [22] defined over $\mathbb{Q}_p$ for torsors, as well as a subvariety $H_g^1(G_v, U^{et})$, consisting of those torsors that trivialize upon base-change to B_{DR} . We warn the reader at this point that this subvariety was denoted H_f^1 in the reference [22]. In this paper, we will use the notation H_f^1 for a certain subvariety of H_g^1 defined using B_{cr} in place of B_{DR} (see the following section). Let us spell out what this B_{DR} condition means. If $T = \text{Spec}(\mathcal{T})$ is a torsor for U^{et} over R , we say it is a *De Rham torsor* if there is a G_v -equivariant algebra homomorphism

$$t : \mathcal{T} \rightarrow R \otimes B_{DR}$$

or equivalently,

$$t_{B_{DR}} : \mathcal{T} \otimes B_{DR} \rightarrow R \otimes B_{DR}.$$

That is, we are requiring the existence of a G_v -invariant point in $T_{B_{DR}}(R \otimes B_{DR})$. Then t induces an isomorphism of B_{DR} -algebras

$$C_t : \mathcal{T} \otimes B_{DR} \simeq A^{et} \otimes R \otimes B_{DR}$$

that is G_v -equivariant. Furthermore, since $t_{B_{DR}}$ is a homomorphism of B_{DR} algebras, it respects the Hodge filtration induced by that on B_{DR} . Hence, so does C_t . Since there is the identity homomorphism

$$e : A^{et} \otimes R \otimes B_{DR} \rightarrow R \otimes B_{DR},$$

out of which we recover t as $e \circ C_t$, we see that the datum of C_t is equivalent to the De Rham condition.

Taking G_v -invariants, we get an isomorphism

$$C_{t^{G_v}} : D(\mathcal{T}) := (\mathcal{T} \otimes B_{DR})^{G_v} \simeq (A^{et} \otimes B_{DR})^{G_v} \otimes R \simeq A^{DR} \otimes R = A^{DR} \otimes_{F_v} (F_v \otimes R)$$

Here, we are using the fact that $D(A^{et}) \simeq A_{DR}$ in a manner compatible with the Hodge filtration [25]. As a consequence of the definition, the coordinate ring $\mathcal{T}$ becomes a limit of finite-dimensional De Rham representations in the sense of [11] and hence, the inclusion

$$D(\mathcal{T}) \hookrightarrow \mathcal{T} \otimes B_{DR}$$

induces an isomorphism

$$D(\mathcal{T}) \otimes_{F_v} B_{DR} \simeq \mathcal{T} \otimes B_{DR}.$$

If we examine what happens to the torsor structure $\mathcal{T} \rightarrow \mathcal{T} \otimes_R (A^{et} \otimes R)$, it extends to

$$\mathcal{T} \otimes B^{DR} \rightarrow \mathcal{T} \otimes_R (A^{et} \otimes R) \otimes B^{DR}$$

But

$$\begin{aligned} \mathcal{T} \otimes_R (A^{et} \otimes R) \otimes B^{DR} &= (\mathcal{T} \otimes A^{et}) \otimes B_{DR} = (\mathcal{T} \otimes B_{DR}) \otimes_{B_{DR}} (A^{et} \otimes B_{DR}) \\ &= ((\mathcal{T} \otimes B_{DR})^{G_v} \otimes_{F_v} B_{DR}) \otimes_{B_{DR}} ((A^{et} \otimes B_{DR})^{G_v} \otimes_{F_v} B_{DR}) = D(\mathcal{T}) \otimes_{F_v} D(A^{et}) \otimes B_{DR} \end{aligned}$$

So taking G_v -invariants gives us

$$D(\mathcal{T}) \rightarrow D(\mathcal{T}) \otimes_{F_v} A^{DR} = D(\mathcal{T}) \otimes_{R \otimes F_v} (A^{DR} \otimes_{F_v} (R \otimes F_v)).$$

This defines the structure of a U^{DR} -torsor over $R \otimes F_v$ on $D(\mathcal{T})$ which is trivialized by the point $t^{G_v} : D(\mathcal{T}) \rightarrow R \otimes F_v$ obtained from t by restricting to the G_v -invariants. In fact, the map

$$D(\mathcal{T}) \rightarrow A^{DR} \otimes_{F_v} (R \otimes F_v)$$

obtained from t^{G_v} is seen to be none other than the earlier isomorphism $C_{t^{G_v}}$. The compatibility of the torsor structure with the Eilenberg-MacLane filtration follows from the fact that the comparison isomorphism of [25] arises from an equivalence of categories

$$\mathrm{Un}_p^{et}(\bar{X}) \otimes B_{DR} \simeq \mathrm{Un}^{DR}(X) \otimes_{F_v} B_{DR}$$

that respects the subcategories specified by the index of unipotency on either side.

The Hodge filtration on $D(\mathcal{T})$ is defined to be the one induced by the Hodge filtration on B_{DR} . Since this is true also of the one on $A^{DR} = D(A^{et})$ and $\mathcal{T} \otimes B_{DR} \simeq A^{et} \otimes R \otimes B_{DR}$ preserves the filtration, so does $D(\mathcal{T}) \simeq A^{DR} \otimes_{F_v} R$. We conclude that the U^{DR} torsor thus obtained can be trivialized together with the Hodge filtration, in the sense of the previous section.

There is a parallel discussion with B_{cr} in place of B_{DR} that defines a *crystalline* U^{et} -torsor T over R to be one that trivializes over B_{cr} or, equivalently, admits a G_v -equivariant isomorphism of torsors

$$\mathcal{T} \otimes B_{cr} \simeq A^{et} \otimes B_{cr}.$$

Using the usual inclusion of B_{cr} into B_{DR} , we see that a crystalline torsor is also De Rham. It is easy to show that the crystalline condition defines a subvariety

$$H_f^1(G_v, U^{et}) \subset H_g^1(G_v, U^{et})$$

However, since the reference [22] dealt explicitly only with H_g , we sketch the modification necessary to represent the crystalline torsors. For each n , consider the functor on $\mathbb{Q}_p$ -algebras

$$H^0(G_v, U_n^{et, B_{cr}} / U_n^{et}) : R \mapsto H^0(G_v, U_n^{et}(B_{cr} \otimes R) / U_n^{et}(R))$$

Then we have

Lemma 5 $H^0(G_v, U_n^{et, B_{cr}} / U_n^{et})$ is representable by an affine variety over $\mathbb{Q}_p$.

The proof is verbatim the same as [22], section 1, proposition 3 and will be therefore omitted. And then, as in the few paragraphs preceding that proposition, we have an exact sequence

$$0 \rightarrow U_n^{et}(R) \rightarrow U_n^{et}(R \otimes B_{cr}) \rightarrow U_n^{et}(R \otimes B_{cr}) / U_n^{et}(R) \rightarrow 0$$

from which we get the connecting homomorphism

$$H^0(G_v, U_n^{et}(R \otimes B_{cr}) / U_n^{et}(R)) \rightarrow H^1(G_v, U_n^{et}(R))$$

which is functorial in R . Thus, it is represented by a map of varieties, and its image is a subvariety of $H^1(U_n^{et}(R))$. But by looking at the crystalline condition as the existence of a G_v -invariant point in $P^{et}(R \otimes B_{cr})$, we see that the image consists exactly of the crystalline torsors (exactly as in the proof of [22], section 2, lemma 6). By passing to the limit over n , we see that the set of isomorphism classes of crystalline torsors for U^{et} is represented by a pro-algebraic subvariety $H_f^1(G_v, U^{et})$ of $H_g^1(G_v, U^{et})$.

If we denote by K the absolutely unramified subfield of F_v , then we get a functor D_{cr} from crystalline U^{et} -torsors over R to (trivial) U^{cr} -torsors over $R \otimes K$ endowed with a Frobenius endomorphism: This functor of course sends $T = \text{Spec}(\mathcal{T})$ to

$$D_{cr}(T) := \text{Spec}((\mathcal{T} \otimes B_{cr})^{G_v}).$$

A discussion entirely parallel to the B_{DR} case shows that $D_{cr}(T)$ is endowed with a Frobenius endomorphism and is a torsor for $U^{cr} = D_{cr}(U^{et})$ in a manner compatible with this extra structure. As always, this torsor is canonically trivial. We have the comparison

$$D(T) \simeq D_{cr}(T) \otimes_K F_v.$$

Thereby, for crystalline torsors T , $D(T)$ is endowed with both a Hodge filtration and a Frobenius.

To summarize, the functor $T \mapsto D(T)$ defines a map from the isomorphism classes of crystalline U^{et} -torsors over R to admissible U^{DR} torsors over $R \otimes F_v$. It is therefore represented by a pro-algebraic map

$$D : H_f^1(G_v, U^{et}) \rightarrow \text{Res}_{\mathbb{Q}_p}^{F_v}(U^{DR}/F^0).$$

Going back from abstract considerations to the study of rational points, each $P^{et}(x)$ defines a class in $H_f^1(G_v, U^{et})$ [25]. We are thus given an étale unipotent Albanese map

$$j_{loc}^{et} : x \mapsto [P^{et}(x)] \in H_f^1(G_v, U^{et}).$$

When we let $L = F$, the number field itself and let $b \in \mathcal{X}(R)$ be an integral point, then the $\Gamma = \text{Gal}(\bar{F}/F)$ -action factors through $G = G_T$ (as in [28], proof of theorem 2.8), with notation following the introduction. As explained in loc. cit., the point here is that there is an exact sequence

$$0 \rightarrow U^{et} \rightarrow U^{et, ar} \rightarrow G_T \rightarrow 0$$

that is split by the base-point b , accounting for the Γ action on U^{et} and also for the twisted actions corresponding to other points. (Here, the middle term is an ‘arithmetic’ fundamental group obtained by pushing out a pro-finite arithmetic fundamental group $\hat{\pi}_1(\mathcal{X}, b)$ using U^{et} .) Therefore, we have a global continuous cohomology pro-variety

$$H^1(G_T, U^{et})$$

and a classifying sub-variety

$$H_f^1(G_T, U^{et}) \subset H^1(G_T, U^{et}),$$

defined to be the inverse image of $H_f^1(G_v, U^{et})$ under the restriction map

$$H^1(G_T, U^{et}) \rightarrow H^1(G_v, U^{et}).$$

Using this environment, we can finally define the global unipotent Albanese map

$$j_{glob}^{et} : x \in \mathcal{X}(R) \mapsto [P^{et}(x)] \in H_f^1(G_T, U^{et}).$$

To conclude, we have constructed a commutative diagram

$$\begin{array}{ccccc}
\mathcal{X}(R) & \rightarrow & \mathcal{X}_v(R_v) & \rightarrow & \text{Res}_{\mathbb{Q}_p}^{F_v}(U^{DR}/F^0)(\mathbb{Q}_p) \\
\downarrow & & \downarrow & \nearrow & \\
H^1(G_T, U^{et}) & \rightarrow & H_f^1(G_v, U^{et})(\mathbb{Q}_p) & &
\end{array}$$

where the commutativity of the triangle on the right is in [10] and [25], following earlier cases of [26], [27].

Passing from here to the finite-level quotients for each n gives us the diagram from the introduction:

$$\begin{array}{ccccc}
\mathcal{X}(R) & \rightarrow & \mathcal{X}_v(R_v) & \rightarrow & \text{Res}_{\mathbb{Q}_p}^{F_v}(U_n^{DR}/F^0)(\mathbb{Q}_p) \\
\downarrow & & \downarrow & \nearrow & \\
H^1(G_T, U_n^{et}) & \rightarrow & H^1(G_v, U_n^{et})(\mathbb{Q}_p) & &
\end{array}$$

3 Comments I

The proof of the statement that conjecture 1 implies the finiteness of $\mathcal{X}(R)$ is exactly as in [22] and in Chabauty's original argument. The dimension assumption implies that for n as in the conjecture, the image of the map

$$H_f^1(G_T, U_n^{et}) \rightarrow U_n^{DR}/F^0$$

is contained in a proper Zariski-closed subset. Hence, so is the image of

$$\mathcal{X}(R) \rightarrow U_n^{DR}/F^0.$$

We deduce that there is a non-zero algebraic function on U_n^{et}/F^0 that vanishes on this image. But by theorem 1, this function is not identically zero on any residue disk of $\mathcal{X}(R_v)$ and is a Coleman function [2]. Therefore, its zero set on $\mathcal{X}(R_v)$ is finite. It is perhaps worth noting that $\mathcal{X}(R_v) \rightarrow U_n^{DR}/F^0$ is at most finite-to-one (this follows from the $n = 2$ case). The essential point we have shown then is that

$$\text{Im}(\mathcal{X}(R)) \subset \text{Im}(\mathcal{X}(R_v)) \cap \text{Im}(H_f^1(G_T, U_n^{et}))$$

and that the latter intersection is finite (of course assuming conjecture 1). Therefore, the strategy of this paper is reminiscent of Serge Lang's suggestion [24] to prove directly that the intersection of a curve with the Mordell-Weil group of its Jacobian is finite, except that the approach here is p-adic and non-abelian.

We proceed to comment on the relation between conjecture 1 and various standard conjectures. Such a connection is based upon a study of various groups of Selmer type. Given a $\mathbb{Q}_p$ -representation V of G_T we will be especially interested in

$$\text{Sel}_T^0(V) := \text{Ker}(H^1(G_T, V) \rightarrow \bigoplus_{w \in T} H^1(G_w, V))$$

Conjecture 2 *Let X/F be a smooth curve and let $V_n = H_{et}^1(\bar{X}, \mathbb{Q}_p)^{\otimes n}(1)$. Then $\text{Sel}_T^0(V_n) = 0$ for $n \gg 0$.*

This conjecture appears to be a special case of some general expectations about motivic representations, and I am not sure to whom it should be attributed. Notice that V_n is a direct summand of $H^n(\bar{X}^n, \mathbb{Q}_p(1))$. There is a Chern class map from K -theory

$$ch_{n,r} : K_{2r-n-1}^{(r)}(X^n) \rightarrow H^1(\text{Gal}(\bar{F}/F), H^n(\bar{X}^n, \mathbb{Q}_p(r)))$$

for $2r - 1 \neq n$ and cycle maps

$$c_r : CH_0^r(X^n) \rightarrow H^1(\text{Gal}(\bar{F}/F), H^{2r-1}(\bar{X}^n, \mathbb{Q}_p(r)))$$

where the superscript in the K -group refers to an associated graded space for the gamma filtration, while the subscript in the Chow group refers to the cycle classes homologically equivalent to zero. The images of $ch_{n,r}$ and c_r lie inside the subspaces $H_g^1(\text{Gal}(\bar{F}/F), H^n(\bar{X}^n, \mathbb{Q}_p(r)))$ consisting of cohomology classes that are unramified at almost all primes and potentially semi-stable at all primes. In particular,

$$\text{Sel}_T^0(V_n) \subset H_g^1(\text{Gal}(\bar{F}/F), H^n(\bar{X}^n, \mathbb{Q}_p(1))).$$

In the course of formulating their famous conjectures, Bloch and Kato conjectured ([3], conjecture 5.3) that $ch_{n,r}$ induces an isomorphism

$$K_{2r-n-1}^{(r)}(X^n) \otimes \mathbb{Q}_p \simeq H_g^1(\text{Gal}(\bar{F}/F), H^n(\bar{X}^n, \mathbb{Q}_p(r)))$$

This can be view as a grand generalization of the finiteness conjecture for Tate-Shafarevich groups of Abelian varieties. It is also stating that certain extensions are geometric in a specific sense. We will explain below how the section conjecture can also be viewed in a similar light. In any case, if $n > 2r - 1$ the K -groups are zero. Therefore

Observation 1 *The conjecture of Bloch and Kato implies conjecture 2.*

Tony Scholl has pointed out to me that

Observation 2 *Conjecture 2 is also implied by the Fontaine-Mazur conjecture.*

This is because an extension

$$0 \rightarrow V_n \rightarrow E \rightarrow \mathbb{Q}_p \rightarrow 0$$

corresponding to an element of this group must be semi-stable at p , and hence, a $\mathbb{Q}_p$ linear combination of extensions of geometric origin. In particular, it must carry a weight filtration. V_n is pure of weight $n - 2$. Since a map between mixed representations is strict for the weight filtration, there exists a vector $v \in W_0 E$ that maps to $1 \in \mathbb{Q}_p$. If $n \geq 3$, then $W_0 E \cap V_n = 0$, and hence, $W_0 E$ must be a 1-dimensional complement to V_n . Thus, the extension must split.

We will show

Proposition 2 *Conjecture 2 implies conjecture 1.*

If we use the symbol U^M for the fundamental group in any of the realizations of interest and $U_n^M = Z^n \backslash U^M$, then we have an exact sequence

$$0 \rightarrow Z^{n+1} \backslash Z^n \rightarrow U_{n+1}^M \rightarrow U_n^M \rightarrow 0$$

We need to calculate the dimension d_n of $Z^{n+1} \backslash Z^n$. This is achieved as follows: U^M is the unipotent completion of either a free group on $m = 2g + s - 1$ generators, where g is the genus of X' and s the order of $X' \setminus X$, or a free group on $2g$ generators modulo a single relation of degree 2 (the compact case). According to [23] d_n is given by the recursive formula

$$\sum_{k|n} k d_k = m^n$$

in the open case and

$$\sum_{k|n} k d_k = (g + \sqrt{g^2 - 1})^n + (g - \sqrt{g^2 - 1})^n$$

in the compact case. So one gets the asymptotics $d_n \approx m^n/n$ in the non-compact case and $d_n \approx (g + \sqrt{g^2 - 1})^n/n$ in the compact case. (The small difference between the two formulas appears to be significant. Even after considerable effort, the argument of observation 3 below could not be adapted to the compact case.) Notice here that in the compact case, $g \geq 2$. We can also estimate $F^0(Z^{n+1} \backslash Z^n)$ in the De Rham realization by noting that it is a quotient of $H_1^{DR}(X_v)^{\otimes n}$ which has the filtration dual to that on $H_{DR}^1(X_v)^{\otimes n}$. So $F^0 H_1^{DR}(X_v)^{\otimes n}$ has dimension equal to the codimension of

$$F^1 H_{DR}^1(X_v)^{\otimes n}$$

$$= [F^1 H_{DR}^1 \otimes (H_{DR}^1)^{\otimes(n-1)}] \oplus [H_{DR}^1 \otimes F^1 H_{DR}^1 \otimes (H_{DR}^1)^{\otimes(n-2)}] \oplus \dots \oplus [(H_{DR}^1)^{\otimes(n-1)} \otimes F^1 H_{DR}^1]$$

which therefore is equal to

$$\dim H^1(\mathcal{O}_{X'_v})^{\otimes n} = g^n.$$

Therefore, $F^0(Z^{n+1} \setminus Z^n) \leq g^n$. In particular, it does not contribute to the asymptotics. That is, the jump in dimensions as one goes from U_n^{DR} to U_{n+1}^{DR} is determined by the asymptotics of d_n . On the other hand, in the étale realization, we have

$$0 \rightarrow H^1(G_T, Z^{n+1} \setminus Z^n) \rightarrow H^1(G_T, U_{n+1}^{et}) \rightarrow H^1(G_T, U_n^{et})$$

in the sense of [22] section 1, whereby the middle term is an $H^1(G_T, Z^{n+1} \setminus Z^n)$ -torsor over a subvariety of the last term. So we can control the change in dimensions using the Euler characteristic formula

$$\dim H^1(G_T, Z^{n+1} \setminus Z^n) - \dim H^2(G_T, Z^{n+1} \setminus Z^n) = \dim(Z^{n+1} \setminus Z^n)^-,$$

where the negative superscript refers to the (-1) eigenspace of complex conjugation. By comparison with complex Hodge theory, we see that the right hand side is $d_n/2$ for n odd. So it remains to observe that

Lemma 6 *Conjecture 2 implies*

$$\dim H^2(G_T, Z^{n+1} \setminus Z^n) \leq P(n)g^n$$

for an effective polynomial $P(n)$ of n .

Proof of lemma. We have a surjection

$$H^2(G_T, H_1(\bar{X}, \mathbb{Q}_p)^{\otimes n}) \rightarrow H^2(G_T, Z^{n+1} \setminus Z^n) \rightarrow 0,$$

so it suffices to prove the estimate for the first group. But by conjecture 2 and Poitou-Tate duality, we get an injection

$$H^2(G_T, H_1(\bar{X}, \mathbb{Q}_p)^{\otimes n}) \hookrightarrow \bigoplus_{w \in T} H^2(G_w, H_1(\bar{X}, \mathbb{Q}_p)^{\otimes n})$$

for n sufficiently large. Hence, we need only estimate the dimension of the local groups. Applying local duality allows us to reduce this to the study of

$$H^0(G_w, H^1(\bar{X}, \mathbb{Q}_p)^{\otimes n}(1)).$$

For w not dividing p , a combination of the geometric weight filtration and the monodromy filtration gives us

$$W : W_0 \subset W_1 \subset W_2 = H^1(\bar{X}, \mathbb{Q}_p)$$

where $\dim Gr_0^W \leq g$. From this, we get immediately that

$$\dim H^0(G_w, H^1(\bar{X}, \mathbb{Q}_p)^{\otimes n}(1)) \leq n \dim Gr_2^W g^{n-1} + \frac{n(n-1)}{2} (\dim Gr_1^W)^2 g^{n-2}$$

For w dividing p , we use the Hodge-Tate decomposition:

$$H^1(\bar{X}, \mathbb{Q}_p) \otimes \mathbb{C}_p \simeq H^0(X', \Omega_{X'}(\log D)) \otimes \mathbb{C}_p(-1) \oplus H^1(X', \mathcal{O}_{X'}) \otimes \mathbb{C}_p$$

as G_w -representations. Thus,

$$H^0(G_w, H^1(\bar{X}, \mathbb{Q}_p)^{\otimes n}(1)) \hookrightarrow H^0(G_w, [H^0(X', \Omega_{X'}(\log D)) \otimes \mathbb{C}_p(-1) \oplus H^1(X', \mathcal{O}_{X'}) \otimes \mathbb{C}_p]^{\otimes n}(1))$$

from which we easily extract the estimate

$$\dim H^0(G_w, H^1(\bar{X}, \mathbb{Q}_p)^{\otimes n}(1)) \leq n(g + s - 1)g^{n-1}$$

□

Since we have already discussed the structure of U^M in the proof of the previous proposition, it becomes easy to outline the relationship to a conjecture of Jannsen. There, one starts with any smooth projective variety V over F having good reduction outside a finite set T of primes (which we take to include all primes dividing p) and considers $H^2(G_T, H^n(\bar{V}, \mathbb{Q}_p)(r))$ with various twists r . Then Jannsen conjectures:

$$H^2(G_T, H^n(\bar{V}, \mathbb{Q}_p)(r)) = 0$$

for $r \geq n + 2$.

In fact, we need only a weaker variant

(Weak Jannsen conjecture) *There exists a $k > 0$ such that for varieties V as above*

$$H^2(G_T, H^n(\bar{V}, \mathbb{Q}_p)(r)) = 0$$

for $r \geq n + k$.

Observation 3 *When X is affine, Jannsen's conjecture in the weaker version implies conjecture 1.*

The point here is to try once more to estimate the H^2 . But in the affine case, the growth rate of $Z^{n+1} \setminus Z^n$ is like m^n/n , so it will certainly suffice to show that $H^2(G_T, H_1(\bar{X}, \mathbb{Q}_p)^{\otimes n})$ is bounded by Ca^n for some $a < m$. Now, easy homological arguments give us the existence of a constant K such that

$$\dim H^2(G_T, V) \leq K \dim V$$

Recall that $m = 2g + s - 1$ and that $H_1(\bar{X}, \mathbb{Q}_p)$ fits into an exact sequence

$$0 \rightarrow W \rightarrow H_1(\bar{X}, \mathbb{Q}_p) \rightarrow V \rightarrow 0$$

where $W = \mathbb{Q}_p(1)^{s-1}$ and $V = H_1(\bar{X}', \mathbb{Q}_p)$. (Here, we have passed to a finite extension where all the points at infinity are rational. This passage is admissible for our H^2 considerations.) Therefore, in the Grothendieck group, we get

$$H_1(\bar{X}, \mathbb{Q}_p)^{\otimes n} = V^{\otimes n} \oplus n(V^{\otimes(n-1)} \otimes W) \oplus \dots \oplus \binom{n}{k}(V^{\otimes(n-k)} \otimes W^k) \oplus \dots$$

But $V \simeq H^1(\bar{X}', \mathbb{Q}_p)(1)$ so that $V^{\otimes l}$ is a direct summand of $H^l((\bar{X}')^l, \mathbb{Q}_p)(l)$. Therefore, $H^2(G_T, V^{\otimes l}(r)) = 0$ for $r \geq k$. Hence,

$$\dim H^2(G_T, H_1(\bar{X}, \mathbb{Q}_p)^{\otimes n}) \leq P(n) \sum_{l=0}^{k-1} \dim H^2(G_T, V^{\otimes(n-l)}(l))$$

for some polynomial $P(n)$. We analyze the cohomology groups on the right. The Weil pairing induces an inclusion

$$\mathbb{Q}_p(1) \hookrightarrow V^{\otimes 2}$$

which continues to an inclusion

$$\mathbb{Q}_p(k) \hookrightarrow V^{\otimes(2k)}$$

with image a direct summand. Write $V^{\otimes 2k} = M \oplus \mathbb{Q}_p(k)$. On the one hand, since any $M^{\otimes i}$ is a direct summand of a $V^{\otimes l}$, we have

$$H^2(G_T, V^{\otimes i} \otimes M^{\otimes j}(r)) = 0$$

for $r \geq k$. On the other,

$$V^{\otimes 2kn} = M^{\otimes n} \oplus M'(k)$$

where M' is a direct sum of (non-negative) tensor products of M and $\mathbb{Q}_p(1)$. Hence,

$$V^{\otimes (2kn+i)} = M^{\otimes n} \otimes V^{\otimes i} \oplus M' \otimes V^{\otimes i}(k)$$

and

$$H^2(G_T, V^{\otimes (2kn+i)}(l)) = H^2(G_T, M^{\otimes n} \otimes V^{\otimes i}(l))$$

for $l \geq 0$. Therefore, as i runs through $0, \dots, 2k-1$, there is a polynomial Q such that

$$\dim H^2(G_T, V^{\otimes (2kn+i)}(l)) \leq Q(n) \dim M^{\otimes n} = Q(n)((2g)^{2k} - 1)^n \leq Q(n)((2g)^{2k} - 1)^{(2kn+i)/(2k)}.$$

Rewriting the exponent, for any $n \geq 0$,

$$\dim H^2(G_T, V^{\otimes n}(l)) \leq Q(n)[((2g)^{2k} - 1)^{1/(2k)}]^n$$

from which the desired estimate follows.

4 Comments II

It is rather interesting to consider the special case of an elliptic curve minus the origin. For example, our method shows finiteness of integral points for CM-elliptic curves E defined over $\mathbb{Q}$ of rank 1, independently of any conjecture. This follows from analyzing somewhat carefully a refined Selmer variety

$$\text{Sel}_T(U_n),$$

defined to be the classes in $H_f^1(G_T, U_n^{et})$ that are potentially unramified at primes not equal to p and potentially crystalline at p . (Here of course, the U refer to the unipotent fundamental groups of E minus the origin.) The definition of this subvariety depends on the construction of certain local subvarieties

$$H_{pf}^1(G_v, H_v; U_n^{et}) \subset H^1(G_v, U_n^{et})$$

$(v|p)$ and

$$H_{pun}^1(G_v, H_v; U_n^{et}) \subset H^1(G_v, U_n^{et})$$

(v not dividing p) for various v and subgroups of finite index $H_v \subset G_v$. Since the pattern is exactly as in [22], section 1, we will omit the details. The point is that if N is a closed subgroup of G , then the restriction map $H^1(G, U_n^{et}) \rightarrow H^1(N, U_n^{et})$ is functorial. We can therefore construct the unramified cohomology varieties $H_{un}^1(G_v, U_n^{et})$ as the inverse image of the base-point in $H^1(I_v, U_n^{et})$, where $I_v \subset G_v$ is the inertia subgroup. For this last object, we need only deal with it when the action of I_v on U_n^{et} is trivial. In this case, the cohomology functor is not necessarily representable by a variety. However, we still have functorial sequences

$$0 \rightarrow H^1(I_v, Z^{n+1} \setminus Z^n) \rightarrow H^1(I_v, U_{n+1}^{et}) \rightarrow H^1(I_v, U_n^{et})$$

which are exact in the naive sense that the inverse image of the base point under the second map is the kernel. (However, the other fibers may be empty.) Now, the condition that a class $c \in H^1(G, U_{n+1}^{et})$ goes to zero in $H^1(I_v, U_{n+1}^{et})$ is an intersection of the condition that it goes to zero in $H^1(I_v, U_n^{et})$, which defines a subvariety by induction, and then the condition that the image in $H^1(I_v, Z^{n+1} \setminus Z^n)$ is trivial. This last space is a vector group, and hence, the resulting condition defines the desired ‘unramified’ subvariety at level $n+1$.

Then, by considering the restriction maps

$$r_v : H^1(G_v, U_n^{et}) \rightarrow H^1(H_v, U_n^{et}),$$

to subgroups of finite index, we get the potentially unramified subvarieties

$$H_{pun}^1(G_v, H_v; U_n^{et}) := r_v^{-1}(H_{un}^1(H_v, U_n^{et}))$$

and potentially crystalline subvarieties

$$H_{pf}^1(G_v, H_v; U_n^{et}) := r_v^{-1}(H_f^1(H_v, U_n^{et})).$$

We know that E acquires good reduction everywhere after base extension to the field F generated by the 12-torsion. We then take the H_v to be the Galois groups of the completions F_v for each place v of F lying above a place of T . Finally, we define

$$H_{pf}^1(G_p, U_n^{et}) := \cap_{v|p} H_{pf}^1(G_p, H_v, U_n^{et}),$$

and

$$H_{pun}^1(G_v, U_n^{et}) := \cap_{w|v} H_{pun}^1(G_v, H_w, U_n^{et}),$$

for $v \neq p$. Then the global group $Sel_T(U_n)$ is defined to be the intersection of the inverse images of all these local groups under the global-to-local restriction maps:

$$Sel_T(U_n) := \cap_{v \in T} [(Res_{loc,v}^{glob})^{-1}(H_{pun/pf}^1(G_v, U_n^{et}))] \subset H^1(G_T, U_n^{et}).$$

With these definitions in place we claim there is an exact sequence

$$0 \rightarrow Sel_T(Z^{n+1} \setminus Z^n) \rightarrow Sel_T(G_T, U_{n+1}^{et}) \rightarrow Sel_T(U_n^{et})$$

again in the sense that the vector group on the left acts freely on the middle with quotient a subvariety of the right hand side. Most of the proof is a consequence of the exactness of

$$0 \rightarrow H^1(G_T, Z^{n+1} \setminus Z^n) \rightarrow H^1(G_T, U_{n+1}^{et}) \rightarrow H^1(G_T, U_n^{et}),$$

(citekim, section 1) leaving us to check that if c_1, c_2 are two elements of $Sel_T(G_T, U_{n+1}^{et})$ mapping to the same class in $Sel_T(G_T, U_n^{et})$, then one is a translate of the other by a class in $Sel_T(Z^{n+1} \setminus Z^n)$. This can be checked locally at each prime. So let w be a prime of F lying above p and let $c_1, c_2 \in Z^1(H_w, U_{n+1}^{et})$ be two cocycles representing classes of $H_f^1(H_w, U_{n+1}^{et})$ and assume they have the same image in $H_f^1(H_w, U_n^{et})$. We already know that $c_2 = c_1 z$ for some cocycle $z \in Z^1(H_w, Z^{n+1} \setminus Z^n)$ and we need to check that it trivializes over B_{cr} . Since both c_1 and c_2 trivialize over B_{cr} , there are elements $u_1, u_2 \in U_{n+1}^{et}(B_{cr})$ such that $c_i(g) = g(u_i^{-1})u_i$ for $g \in H_w$. Then $z(g) = g(u_1 u_2^{-1})u_2 u_1^{-1}$. Therefore, we see that $u_2 u_1^{-1}$ maps to an H_w -invariant element in $U_n^{et}(B_{cr})$. But the invariant points just comprise U_n^{DR} and the map $U_{n+1}^{DR} \rightarrow U_n^{DR}$ is surjective. Therefore, there is an H_w -invariant element $v \in U_{n+1}(B_{cr})$ such that $u_2 u_1^{-1} = vc$ for some $c \in Z^{n+1} \setminus Z^n(B_{cr})$. Hence, $z(g) = g((vc)^{-1})(vc) = g(c^{-1})c$, giving us the triviality we want. (Of course we should have included a $\mathbb{Q}_p$ -algebra R in the notation to prove all this functorially, but we have omitted it for brevity.) At a prime w not dividing p , the condition to check concerns unramified cocycles, and hence is easier: Suppose c_1, c_2 are two unramified cocycles with values in U_{n+1}^{et} that map to the same class in $H^1(H_w, U_n^{et})$. Then $c_2 = c_1 z$ for some cocycle z with values in $Z^{n+1} \setminus Z^n$. But since the action on U_{n+1}^{et} itself is unramified, all cocycles restricted to the inertia group I_w are homomorphisms. Then since c_1 and c_2 are in fact trivial on I_w , so is z .

For $n = 2$, we get from this the inequality of dimensions

$$\dim Sel_T(U_3^{et}) \leq \dim Sel_T(U_2^{et}) + \dim Sel_T(Z^3 \setminus Z^2).$$

However,

$$Z^3 \setminus Z^2 \simeq H_1(\bar{E}, \mathbb{Q}_p) \wedge H_1(\bar{E}, \mathbb{Q}_p) \simeq \mathbb{Q}_p(1).$$

This implies that $\text{Sel}_T(Z^3 \setminus Z^2)$ consists of the units in $\mathbb{Z}$ tensor $\mathbb{Q}_p$, and hence, is zero ([3], example 3.9). Therefore,

$$\dim \text{Sel}_T(U_3^{et}) = \dim \text{Sel}_T(U_2^{et}) = \dim \text{Sel}_T(H_1(\bar{E}, \mathbb{Q}_p)) = 1,$$

by the rank one hypothesis.

On the De Rham side, we have the same dimension count ($= 1$) for $Z^3 \setminus Z^2$. Meanwhile, considering the quotient map

$$H_1^{DR}(E_p) \otimes H_1^{DR}(E_p) \rightarrow Z^3 \setminus Z^2$$

and the fact that $F^0 H_1 \wedge F^0 H_1 = 0$, we get $F^0(Z^3 \setminus Z^2) = 0$. So

$$\dim F^0 U_3^{DR} = \dim F^0 U_2^{DR} = 1$$

and

$$\dim U_3^{DR}/F^0 = 3 - 1 = 2.$$

Therefore, j_3^{DR} suffices to finish the job.

This calculation shows that in explicit situations like this with a fixed n , it is worth imposing extra Selmer conditions to improve the dimension estimate, in contrast to the general finiteness problem where refined conditions appear to be irrelevant. It is probably also worth looking into this case more carefully with a view towards working out explicit bounds in the manner of Coleman's refinement of Chabauty [7]. The suspicion is that this will require understanding the Dieudonné map D more precisely, along the lines of 'non-abelian explicit reciprocity laws' à la Kato [21].

It was pointed out to me by John Coates that the general CM case is intimately related to a pseudo-nullity conjecture from the Iwasawa theory of elliptic curves. That is, let F_∞ be the field generated by the p -power torsion of E , F_1 the field generated by the p -torsion, $G_\infty = \text{Gal}(F_\infty/F_1)$, and $\Lambda = \mathbb{Z}_p[[G_\infty]]$ the Iwasawa algebra of G_∞ . Let I be the Galois group of the maximal abelian unramified pro- p extension of F_∞ split over the primes lying above T . Coates and Sujatha have conjectured [8] that I is pseudo-null as a Λ -module.

Observation 4 *This pseudo-nullity implies conjecture 1 for $E \setminus \{0\}$.*

Recall that what is necessary is to estimate $H^2(G_T, Z^{n+1} \setminus Z^n)$ which is implied by an estimate for $H^2(G_T, H_1(\bar{E}, \mathbb{Q}_p)^{\otimes n})$. But for just this part, we can assume that the base-field is F_1 since the corestriction map is surjective on H^2 . In fact, the pseudo-nullity implies that

$$\dim H^2(G_T, H_1(\bar{E}, \mathbb{Q}_p)^{\otimes n}) \leq P(n)$$

for some polynomial $P(n)$. To see this, first note that it suffices to prove such an inequality for the Selmer group

$$\text{Ker}(H^2(G_T, H_1(\bar{E}, \mathbb{Q}_p)^{\otimes n}) \rightarrow \oplus_{v \in T} H^2(G_v, H_1(\bar{E}, \mathbb{Q}_p)^{\otimes n}))$$

since the local contributions can be bounded exactly as in section 3 (note that $g = 1$ in this case). So it suffices to bound the H^1 Selmer group

$$\text{Sel}_T^0(H^1(\bar{E}, \mathbb{Q}_p)^{\otimes n}(1)) = \text{Ker}(H^1(G_T, H^1(\bar{E}, \mathbb{Q}_p)^{\otimes n}(1)) \rightarrow \oplus_{v \in T} H^1(G_v, H^1(\bar{E}, \mathbb{Q}_p)^{\otimes n}(1))),$$

and we have an isomorphism:

$$\text{Sel}_T^0(H^1(\bar{E}, \mathbb{Q}_p)^{\otimes n}(1)) \simeq \text{Hom}_\Lambda(I, H^1(\bar{E}, \mathbb{Q}_p)^{\otimes n}(1))$$

coming from the Hochschild-Serre spectral sequence. To see this, it is perhaps convenient to utilize the language of continuous étale cohomology [20] applied to the inverse system

$$(T_i) = (H^1(\bar{E}, \mathbb{Z}/p^i)^{\otimes n}(1)).$$

The Hochschild-Serre spectral sequence in this context gives us the exact sequence

$$0 \rightarrow H^1(G_\infty, (H^0(N, T_i))) \rightarrow H^1(G_T, (T_i)) \rightarrow H^0(G_\infty, (H^1(N, T_i))) \rightarrow H^2(G_\infty, (H^0(N, T_i)))$$

([20], theorem 3.3), where $N \subset G_T$ is the subgroup fixing F_∞ . Since the N -action is trivial, all the $H^0(N, T_i)$ are just T_i , and hence, satisfy the Mittag-Leffler condition. Therefore, all the terms but one become readily identified with the continuous Galois cohomology of $T = \varprojlim T_i = H^1(\bar{E}, \mathbb{Z}_p)^{\otimes n}(1)$ while

$$H^0(G_\infty, (H^1(N, T_i))) := \varprojlim \text{Hom}_{G_\infty}(N, T_i) = \text{Hom}_{G_\infty}(N, T).$$

When we tensor with $\mathbb{Q}$, we get therefore

$$H^1(G_T, H^1(\bar{E}, \mathbb{Q}_p)^{\otimes n}(1)) \simeq \text{Hom}_\Lambda(N, H^1(\bar{E}, \mathbb{Q}_p)^{\otimes n}(1)).$$

Passing to the Selmer group, the last group gets replaced by $\text{Hom}_\Lambda(I, H^1(\bar{E}, \mathbb{Q}_p)^{\otimes n}(1))$.

But the pseudo-nullity implies that I is finitely generated over $\mathbb{Z}_p$ (see, e.g., [14], section 2, paragraph following lemma 2). Thus, if we fix a Frobenius element F in G_∞ , then only finitely many eigenvalues can occur for F in $I \otimes \mathbb{Q}_p$. Weight considerations imply then that

$$\text{Hom}_\Lambda(I, H^1(\bar{E}, \mathbb{Q}_p)^{\otimes n}(1)) = 0$$

for $n \gg 0$.

5 Comments III

Let us consider the section conjecture briefly in terms of Galois cohomology in order to relate it to the Selmer varieties that occur in this paper. For simplicity, we consider it in the situation where a point b exists in $X(F)$ and is used as the base point for the fundamental group. So we have the exact sequence

$$0 \rightarrow \hat{\pi}_1(\bar{X}, b) \rightarrow \hat{\pi}_1(X, b) \rightarrow \Gamma \rightarrow 0$$

of profinite fundamental groups from the introduction as well as one fixed splitting

$$s : \Gamma \rightarrow \hat{\pi}_1(X, b).$$

We use this splitting to make Γ act on $\hat{\pi}_1(\bar{X}, b)$ by conjugation via s . Suppose $t : \Gamma \rightarrow \hat{\pi}_1(X, b)$ is another splitting. Then for each $g \in \Gamma$, there is a $z_t(g) \in \pi_1(\bar{X}, b)$ such that $t(g) = z_t(g)s(g)$. We have $t(g)t(h) = t(gh) = z_t(gh)s(gh)$ and hence, $z_t(gh)s(g)s(h) = z(g)s(g)z_t(h)s(h)$, or $z_t(gh) = z_t(g)g(z_t(h))$. That is to say, z_t is a 1-cocycle. In fact, a straightforward computation shows that this map sets up a bijection

$$[Split(X)/\sim] \simeq H^1(\Gamma, \hat{\pi}_1(\bar{X}, b))$$

to a continuous cohomology set. In the case of a point $x \in X(F)$, the corresponding cohomology class is the class of the torsor $\hat{\pi}_1(\bar{X}; x, b)$ of étale paths from b to x . So we have simply reformulated the section conjecture in terms of a classifying map

$$X(F) \rightarrow H^1(\Gamma, \hat{\pi}_1(\bar{X}, b))$$

that goes to a classifying space for torsors similar to the ones occurring in this paper. In fact, this map is easily seen to be injective so that the content of the section conjecture lies in surjectivity. That is, it is saying that all torsors are of ‘geometric origin’ in a very precise sense.

Let us try to formulate the correct analogue of the section conjecture for an elliptic curve E equipped with the origin $o \in E$ (which we are *not* deleting). We still have an exact sequence

$$0 \rightarrow \hat{\pi}_1(\bar{E}, o) \rightarrow \hat{\pi}_1(E, o) \rightarrow \Gamma \rightarrow 0$$

and the splittings are still classified by $H^1(G, \hat{\pi}_1(\bar{E}, o))$. However, in this non-hyperbolic situation, the splittings that come from points satisfy local conditions that amount to exactly the subspace

$$H_g^1(G, \hat{\pi}_1(\bar{E}, o)) \subset H^1(G, \hat{\pi}_1(\bar{E}, o))$$

studied by Bloch-Kato. That is, the correct classifying map to study in this case is

$$E(F) \rightarrow H_g^1(G, \hat{\pi}_1(\bar{E}, o)).$$

Since

$$\hat{\pi}_1(\bar{E}, o) = H_1(\bar{E}, \hat{\mathbb{Z}}) \simeq H^1(\bar{E}, \hat{\mathbb{Z}})(1),$$

we conclude that the elliptic curve analogue of the section conjecture is nothing but the finiteness conjecture for the Shafarevich group of E . Of course the map that occurs here is identified with the cycle map coming from $CH_0^1(E) \simeq E(F)$. In this abelian situation, the Chern class maps from higher K -theory appear not to play a role in the ‘section conjecture.’

In view of the analogy just outlined, it is perhaps reasonable to think of the relevant parts of the Bloch-Kato conjecture in the hyperbolic case as comprising a ‘linear analogue’ of the section conjecture. In other words, one possible perspective is that we have proved an implication of the form

‘linear’ section conjecture $\Rightarrow$ Faltings’ theorem.

Within this restricted context, one can consider the Fontaine-Mazur conjecture as being a ‘weak’ Bloch-Kato/BSD conjecture, in that one is merely asking that certain extensions be geometric in whatever manner, as opposed to the precise manner of coming from points or the K -theory of specific varieties under discussion. Therefore, it seems natural to ask for a weak section conjecture, whereby one demands that every class in $H^1(\Gamma, \hat{\pi}_1(\bar{X}, b))$ be of geometric origin in some definite sense that lacks as yet a good formulation. (For example, coming to grips with such a formulation appears to be considerably more difficult than a well-known notion from the theory of mixed motives: ‘a system of realizations of geometric origin.’) Meanwhile, for these vague ideas to acquire more weight, it would be nice to prove an implication of the form

section conjecture $\Rightarrow$ linear section conjecture.

Acknowledgements:

-The author was supported in part by a grant from the National Science Foundation.

Deep gratitude is due to:

- Shinichi Mochizuki for overarching influence stemming from his ‘non-linear’ philosophy;
- Dick Hain for continuing to provide patient answers to questions on Hodge theory;
- Bill McCallum, for emphasizing over the years the importance of Chabauty’s method;
- Dinesh Thakur, for a careful and enthusiastic reading of the manuscript;
- Martin Olsson, for communicating to me his deep results on non-abelian p -adic Hodge theory;
- Uwe Jannsen, Guido Kings, Jan Nekovar, and Tony Scholl, for helpful discussions on mixed motives;
- Gerd Faltings, Florian Pop, John Coates, and Kazuya Kato for persistent encouragement of this research and many enlightening discussions on fundamental groups and Iwasawa theory;
- and, of course, Serge Lang, for too many things to be listed.

References

- [1] Berthelot, Pierre Finitude et pureté cohomologique en cohomologie rigide. With an appendix in English by Aise Johan de Jong. Invent. Math. 128 (1997), no. 2, 329–377.
- [2] Besser, Amnon Coleman integration using the Tannakian formalism. Math. Ann. 322 (2002), no. 1, 19–48.

- [3] Bloch, Spencer; Kato, Kazuya L-functions and Tamagawa numbers of motives. *The Grothendieck Festschrift, Vol. 1*, 333–400, Prog. Math. 86, Birkhäuser, Boston, MA 1990.
- [4] Chabauty, Claude Sur les points rationnels des courbes algébriques de genre supérieur l'unité. C. R. Acad. Sci. Paris 212, (1941). 882–885.
- [5] Chen, Kuo-Tsai Iterated path integrals. Bull. Amer. Math. Soc. 83 (1977), no. 5, 831–879.
- [6] Chiarellotto, Bruno; Le Stum, Bernard F -isocristaux unipotents. Compositio Math. 116 (1999), no. 1, 81–110.
- [7] Coleman, Robert F. Effective Chabauty. Duke Math. J. 52 (1985), no. 3, 765–770.
- [8] Coates, J.; Sujatha, R. Fine Selmer groups of elliptic curves over p -adic Lie extensions. Math. Ann. 331 (2005), no. 4, 809–839.
- [9] Deligne, Pierre Le groupe fondamental de la droite projective moins trois points. Galois groups over $\mathbb{Q}$ (Berkeley, CA, 1987), 79–297, Math. Sci. Res. Inst. Publ., 16, Springer, New York, 1989.
- [10] Faltings, Gerd Mathematics around Kim's new proof of Siegel's theorem. Preprint to be published in the proceedings of di Giorgi conference on number theory.
- [11] Fontaine, Jean-Marc Sur certains types de représentations p -adiques du groupe de Galois d'un corps local; construction d'un anneau de Barsotti-Tate. Ann. of Math. (2) 115 (1982), no. 3, 529–577.
- [12] Fontaine, Jean-Marc; Mazur, Barry Geometric Galois representations. Elliptic curves, modular forms, & Fermat's last theorem (Hong Kong, 1993), 41–78, Ser. Number Theory, I, Internat. Press, Cambridge, MA, 1995.
- [13] Furusho, Hidekazu p -adic multiple zeta values I – p -adic multiple polylogarithms and the p -adic KZ equation. Invent. Math. 155 (2004), no. 2, 253–286.
- [14] Greenberg, Ralph On the structure of certain Galois groups. Invent. Math. 47 (1978), no. 1, 85–99.
- [15] Grothendieck, Alexandre Brief an G. Faltings, *Geometric Galois Actions, 1*, 49–58, London Math. Soc. Lecture Note Ser., 242, Cambridge University Press, Cambridge, 1997.
- [16] Hain, Richard M. Higher Albanese manifolds. Hodge theory (Sant Cugat, 1985), 84–91, Lecture Notes in Math., 1246, Springer, Berlin, 1987.
- [17] Hain, Richard M. The de Rham homotopy theory of complex algebraic varieties. I. K -Theory 1 (1987), no. 3, 271–324.
- [18] Hain, Richard M.; Zucker, Steven Unipotent variations of mixed Hodge structure. Invent. Math. 88 (1987), no. 1, 83–124.
- [19] Jannsen, Uwe On the l -adic cohomology of varieties over number fields and its Galois cohomology. Galois groups over $\mathbb{Q}$ (Berkeley, CA, 1987), 315–360, Math. Sci. Res. Inst. Publ., 16, Springer, New York, 1989.
- [20] Jannsen, Uwe Continuous étale cohomology. Math. Ann. 280 (1988), no. 2, 207–245.
- [21] Kato, Kazuya p -adic Hodge theory and values of zeta functions of modular forms. Cohomologies p -adiques et applications arithmétiques. III. Astrisque No. 295 (2004), ix, 117–290.
- [22] Kim, Minhyong The motivic fundamental group of $\mathbf{P}^1 \setminus \{0, 1, \infty\}$ and the theorem of Siegel. Invent. Math. 161 (2005), no. 3, 629–656.

- [23] Labute, John P. On the descending central series of groups with a single defining relation. *J. Algebra* 14 1970 16–23.
- [24] Lang, Serge Division points on curves. *Ann. Mat. Pura Appl.* (4) 70 1965 229–234.
- [25] Olsson, Martin Towards non-abelian p -adic Hodge theory in the good reduction case, preprint. <http://www.math.utexas.edu/~molsson>. (Revised version, in preparation, to comprise the Hodge filtration.)
- [26] Shiho, Atsushi Crystalline fundamental groups and p -adic Hodge theory. The arithmetic and geometry of algebraic cycles (Banff, AB, 1998), 381–398, CRM Proc. Lecture Notes, 24, Amer. Math. Soc., Providence, RI, 2000.
- [27] Vologodsky, Vadim Hodge structure on the fundamental group and its application to p -adic integration. *Mosc. Math. J.* 3 (2003), no. 1, 205–247, 260.
- [28] Wildeshaus, Jörg Realizations of polylogarithms. *Lecture Notes in Mathematics*, 1650. Springer-Verlag, Berlin, 1997.
- [29] Wojtkowiak, Zdzisław Cosimplicial objects in algebraic geometry. Algebraic K -theory and algebraic topology (Lake Louise, AB, 1991), 287–327, NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci., 407, Kluwer Acad. Publ., Dordrecht, 1993.

DEPARTMENT OF MATHEMATICS, PURDUE UNIVERSITY, WEST LAFAYETTE, INDIANA 47907 and DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ARIZONA, TUCSON, AZ 85721, U.S.A.

EMAIL: kimm@math.purdue.edu